

B2

CORRECTED VERSION

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
18 January 2001 (18.01.2001)

PCT

(10) International Publication Number  
WO 01/004322 A1

(51) International Patent Classification<sup>7</sup>: C12N 15/53,  
9/04, C12P 13/08, C12N 1/21, C12Q 1/68 // (C12N 1/21,  
C12R 1:15) (C12P 13/08, C12R 1:15)

View, Roscrea, County Tipperary (IE). BURKE, Kevin;  
5, Greenfields Road, Newcastle, Galway, County Galway  
(IE). MORITZ, Bernd; Schattbachstrasse 1, D-44801  
Bochum (DE). EGGELE, Lothar; Elsenkamp 6,  
D-52428 Jülich (DE). SAHM, Hermann; Wendeli-  
nusstrasse 71, D-52428 Jülich (DE). MÖCKEL, Bettina;  
Benrodestrasse 35, D-40597 Düsseldorf (DE). WEIS-  
SENBORN, Anke; Falkenweg 66, D-72076 Tübingen  
(DE).

(21) International Application Number: PCT/EP00/06300

(22) International Filing Date: 5 July 2000 (05.07.2000)

(25) Filing Language: English

(26) Publication Language: English

(81) Designated States (national): AU, BR, CA, CN, HU, ID,  
JP, KR, MX, PL, RU, SK, UA, ZA.

(30) Priority Data:  
60/142,915 9 July 1999 (09.07.1999) US  
09/531,267 20 March 2000 (20.03.2000) US

(84) Designated States (regional): European patent (AT, BE,  
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,  
NL, PT, SE).

(71) Applicant: DEGUSSA AG [DE/DE]; Bennigsenplatz 1,  
40474 Düsseldorf (DE).

Published:  
with international search report

(71) Applicants (for AT, AU, BE, BR, CA, CH, CN, CY, DE,  
ES, FI, FR, GB, GR, HU, ID, IT, JP, KR, LU, MC, MX,  
NL, PT, RU, SE, SK, UA, ZA only): FORSCHUNGSZEN-  
TRUM JÜLICH GMBH [DE/DE]; D-52425 Jülich (DE).  
NATIONAL UNIVERSITY OF IRELAND [IE/IE]; Gal-  
way (IE).

(48) Date of publication of this corrected version:  
12 September 2002

(15) Information about Correction:  
see PCT Gazette No. 37/2002 of 12 September 2002, Sec-  
tion II

(72) Inventors: DUNICAN, L., K. (deceased). MC-  
CORMACK, Ashling; Moate Road, Athlone, County  
Westmeath (IE). STAPELTON, Cliona; 27, Railway

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

WO 01/004322 A1

(54) Title: NUCLEOTIDE SEQUENCES WHICH CODE FOR THE OPCA GENE

(57) Abstract: The invention relates to an isolated polynucleotide from coryneform bacteria, comprising at least one polynucleotide sequence chosen from the group consisting of a) polynucleotide which is identical to the extent of at least 70 % to a polynucleotide which codes for polypeptides which comprise at least one of the amino acid sequences according to SEQ ID No. 3 or SEQ ID No. 5 or SEQ ID No. 8 or SEQ ID No. 10, b) polynucleotide which codes for polypeptides which comprise amino acid sequences which are identical to the extent of at least 70 % to the amino acid sequences according to SEQ ID No. 3 or SEQ ID No. 5 or according to SEQ ID No. 8 or SEQ ID No. 10, c) polynucleotide which is complementary to the polynucleotides of a) or b), or d) polynucleotide comprising at least 15 successive nucleotides of the polynucleotide sequences of a), b) or c), and a process for the fermentative preparation of L-amino acids, which comprises in the coryneform microorganisms which in particular already produce L-amino acids, a) amplifying, in particular over-expressing, in addition to the opcA gene, at least one of the nucleotide sequences which code for the tal gene, tkt gene, zwf gene or the devB gene. b) concentrating the desired L-amino acid in the medium or in the cells of the bacteria and c) isolating the L-amino acid.

### New nucleotide sequences which code for the opcA gene

The invention provides nucleotide sequences which code for the opcA gene and a process for the fermentative preparation of amino acids, in particular L-lysine using  
5 coryneform bacteria in which the opcA gene is amplified.

#### Prior art

Amino acids, in particular L-lysine, are used in human medicine and in the pharmaceuticals industry, but in particular in animal nutrition.

10 It is known that amino acids are prepared by fermentation from strains of coryneform bacteria, in particular *Corynebacterium glutamicum*. Because of their great importance, work is constantly being undertaken to improve the preparation processes. Improvements to the processes  
15 can relate to fermentation measures, such as e. g. stirring and supply of oxygen, or the composition of the nutrient media, such as e. g. the sugar concentration during the fermentation, or the working up to the product form by e. g. ion exchange chromatography, or the intrinsic output  
20 properties of the microorganism itself.

Methods of mutagenesis, selection and mutant selection are used to improve the output properties of these microorganisms. Strains which are resistant to antimetabolites, such as e. g. the lysine analogue S-(2-aminoethyl)-cysteine, or are auxotrophic for metabolites of  
25 regulatory importance and produce L-amino acids, such as e. g. L-lysine, are obtained in this manner. Methods of the recombinant DNA technique have also been employed for some years for improving the strain of *Corynebacterium* strains  
30 which produce amino acids.

The importance of the pentose phosphate cycle for the biosynthesis is known.

- Thus Oishi and Aida (Agricultural and Biological Chemistry 29, 83-89 (1965)) already report on the "hexose monophosphate shunt" of *Brevibacterium ammoniagenes*. Sugimoto and Shio (Agricultural and Biological Chemistry 51, 101-108 (1987)) report on the regulation of glucose 6-phosphate dehydrogenase in *Brevibacterium flavum*. Sugimoto and Shio (Agricultural and Biological Chemistry 51, 1257-11263 (1987)) report on the regulation of glucose 6-phosphate dehydrogenase in *Brevibacterium flavum*.
- 10 JP-A-09224661 discloses the nucleotide sequence of the glucose 6-phosphate dehydrogenase gene, called *zwf*, of *Brevibacterium flavum* MJ-223 (FERM BP-1497). JP-A-09224661 describes the N-terminal amino acid sequence of the *Zwf* polypeptide as Met Val Ile Phe Gly Val Thr Gly Asp Leu Ala
- 15 Arg Lys Lys Leu.

However, it has not been possible to confirm this.

#### Object of the invention

- The inventors had the object of providing new measures for improved fermentative preparation of amino acids, in particular L-lysine.
- 20

#### Description of the invention

- Amino acids, in particular L-lysine, are used in human medicine, in the pharmaceuticals industry and in particular in animal nutrition. There is therefore a general interest in providing new improved processes for the preparation of amino acids, in particular L-lysine.
- 25

- When L-lysine or lysine are mentioned in the following, not only the base but also the salts, such as e. g. lysine monohydrochloride or lysine sulfate, are also meant by this.
- 30

The invention provides an isolated polynucleotide from coryneform bacteria, comprising at least one polynucleotide sequence chosen from the group consisting of

- 5 a) polynucleotide which is identical to the extent of at least 70 % to a polynucleotide which codes for polypeptides which comprise at least one of the amino acid sequences according to SEQ ID No. 3 or SEQ ID No. 5 or SEQ ID No. 8 or SEQ ID No. 10,
- 10 b) polynucleotide which codes for polypeptides which comprise amino acid sequences which are identical to the extent of at least 70 % to the amino acid sequences according to SEQ ID No. 3 or SEQ ID No. 5 or according to SEQ ID No. 8 or SEQ ID No. 10,
- 15 c) polynucleotide which is complementary to the polynucleotides of a) or b), or
- d) polynucleotide comprising at least 15 successive nucleotides of the polynucleotide sequences of a), b) or c).

20 The invention also provides the polynucleotide as claimed in claim 1, this preferably being a DNA which is capable of replication, comprising:

- (i) one or more nucleotide sequence(s) chosen from the group consisting of SEQ ID No. 1, SEQ ID No. 4, SEQ ID No. 6, SEQ ID No. 9, or
- 25 (ii) at least one sequence which corresponds to sequence (i) within the range of the degeneration of the genetic code, or
- (iii) at least one sequence which hybridizes with the sequence complementary to sequence (i) or (ii),
- 30 and optionally
- (iv) sense mutations of neutral function in (i).

The invention also provides

a polynucleotide as claimed in claim 4, comprising the nucleotide sequence as shown in SEQ ID No. 4 or SEQ ID No. 9,

- 5 a polynucleotide as claimed in claim 6, which codes for a polypeptide which comprises at least one of the amino acid sequences as shown in SEQ ID No. 3, SEQ ID No. 5, SEQ ID No. 8 or SEQ ID No. 10,

- a vector containing the polynucleotide as claimed in  
10 claim 1,

and coryneform bacteria, serving as the host cell, which contain the vector.

- The invention also provides polynucleotides which substantially comprise a polynucleotide sequence, which are  
15 obtainable by screening by means of hybridization of a corresponding gene library, which comprises the complete gene with the polynucleotide sequence corresponding to SEQ ID No. 4 or SEQ ID No. 9, with a probe which comprises the sequence of the polynucleotide mentioned, according to SEQ  
20 ID No. 4 or SEQ ID No. 9 or a fragment thereof, and isolation of the DNA sequence mentioned.

- Polynucleotide sequences according to the invention are suitable as hybridization probes for RNA, cDNA and DNA, in order to isolate, in the full length, cDNA which code for  
25 Opca protein and to isolate those cDNA or genes which have a high similarity of sequence with that of the opca gene.

- Polynucleotide sequences according to the invention are furthermore suitable as primers for the preparation of DNA of genes which code for Opca protein by the polymerase  
30 chain reaction (PCR).

Such oligonucleotides which serve as probes or primers comprise at least 30, preferably at least 20, especially preferably at least 15 successive nucleotides. Oligonucleotides which have a length of at least 40 or 50 nucleotides are also suitable.

"Isolated" means separated out of its natural environment.

"Polynucleotide" in general relates to polyribonucleotides and polydeoxyribonucleotides, it being possible for these to be non-modified RNA or DNA or modified RNA or DNA.

10 "Polypeptides" is understood as meaning peptides or proteins which comprise two or more amino acids bonded via peptide bonds.

The polypeptides according to the invention include a polypeptide according to SEQ ID No. 3, SEQ ID No. 5, SEQ ID  
15 No. 8 or SEQ ID No. 10, in particular those with the biological activity of the OpcA gene product, and also those which are identical to the extent of at least 70 % to the polypeptide according to SEQ ID No. 3, SEQ ID No. 5, SEQ ID No. 8 or SEQ ID No. 10, and preferably are identical  
20 to the extent of at least 80% and in particular to the extent of at least 90 % to 95 % to the polypeptide according to SEQ ID No. 3, SEQ ID No. 5, SEQ ID No. 8 or SEQ ID No. 10, and have the activity mentioned.

The invention also provides the new Zwf protein which forms  
25 the Zwf sub-unit of glucose 6-phosphate dehydrogenase. The amino acid sequence of the translation product is shown in SEQ ID no. 2 and SEQ ID No. 7. The N-terminal amino acid sequence of the Zwf sub-unit, which can be isolated, of glucose 6-phosphate dehydrogenase is shown in SEQ ID No.  
30 11.

The invention also provides a process for the fermentative preparation of amino acids, in particular L-lysine, using coryneform bacteria which in particular already produce an

amino acid, and in which the nucleotide sequences which code for the *opcA* gene are amplified, in particular over-expressed, optionally together with the *zwf* gene.

The term "amplification" in this connection describes the increase in the intracellular activity of one or more enzymes (proteins) in a microorganism which are coded by the corresponding DNA, for example by increasing the number of copies of the gene or genes, using a potent promoter or using a gene which codes for a corresponding enzyme (protein) having a high activity, and optionally combining these measures.

The microorganisms which the present invention provides can prepare L-amino acids, in particular L-lysine, from glucose, sucrose, lactose, fructose, maltose, molasses, starch, cellulose or from glycerol and ethanol. They can be representatives of coryneform bacteria, in particular of the genus *Corynebacterium*. Of the genus *Corynebacterium*, there may be mentioned in particular the species *Corynebacterium glutamicum*, which is known among experts for its ability to produce L-amino acids.

Suitable strains of the genus *Corynebacterium*, in particular of the species *Corynebacterium glutamicum*, are, for example, the known wild-type strains

*Corynebacterium glutamicum* ATCC13032  
*Corynebacterium acetoglutamicum* ATCC15806  
*Corynebacterium acetoacidophilum* ATCC13870  
*Corynebacterium thermoaminogenes* FERM BP-1539  
*Corynebacterium melassecola* ATCC17965  
*Brevibacterium flavum* ATCC14067  
*Brevibacterium lactofermentum* ATCC13869 and  
*Brevibacterium divaricatum* ATCC14020

and L-lysine-producing mutants or strains prepared therefrom, such as, for example

Corynebacterium glutamicum FERM-P 1709  
Brevibacterium flavum FERM-P 1708  
Brevibacterium lactofermentum FERM-P 1712  
Corynebacterium glutamicum FERM-P 6463  
5 Corynebacterium glutamicum FERM-P 6464 and  
Corynebacterium glutamicum DSM5715  
Corynebacterium glutamicum DM58-1  
Corynebacterium glutamicum DSM12866.

and L-threonine-producing mutants or strains prepared  
10 therefrom, such as, for example

Corynebacterium glutamicum ATCC21649  
Brevibacterium flavum BB69  
Brevibacterium flavum DSM5399  
Brevibacterium lactofermentum FERM-BP 269  
15 Brevibacterium lactofermentum TBB-10

and L-isoleucine-producing mutants or strains prepared  
therefrom, such as, for example

Corynebacterium glutamicum ATCC 14309  
Corynebacterium glutamicum ATCC 14310  
20 Corynebacterium glutamicum ATCC 14311  
Corynebacterium glutamicum ATCC 15168  
Corynebacterium ammoniagenes ATCC 6871

and L-tryptophan-producing mutants or strains prepared  
therefrom, such as, for example

25 Corynebacterium glutamicum ATCC21850 and  
Corynebacterium glutamicum KY9218 (pKW9901)

The inventors have succeeded in isolating the new *opcA* gene  
of *C. glutamicum* which codes for the *OpcA* sub-unit of the  
enzyme glucose 6-phosphate dehydrogenase (EC 2.7.1.11).

30 To isolate the *opcA* gene or also other genes of *C.*  
*glutamicum*, a gene library of this microorganism is first



set up in *E. coli*. The setting up of gene libraries is described in generally known textbooks and handbooks. The textbook by Winnacker: *Gene und Klone, Eine Einführung in die Gentechnologie* [Genes and Clones, An Introduction to Genetic Engineering] (Verlag Chemie, Weinheim, Germany, 1990) or the handbook by Sambrook et al.: *Molecular Cloning, A Laboratory Manual* (Cold Spring Harbor Laboratory Press, 1989) may be mentioned as an example. A very well-known gene library is that of the *E. coli* K-12 strain W3110 set up in  $\lambda$  vectors by Kohara et al. (Cell 50, 495-508 (1987)). Bathe et al. (Molecular and General Genetics, 252:255-265, 1996) describe a gene library of *C. glutamicum* ATCC13032, which was set up with the aid of the cosmid vector SuperCos I (Wahl et al., 1987, Proceedings of the National Academy of Sciences USA, 84:2160-2164) in the *E. coli* K-12 strain NM554 (Raleigh et al., 1988, Nucleic Acids Research 16:1563-1575). Börmann et al. (Molecular Microbiology 6(3), 317-326) (1992) in turn describe a gene library of *C. glutamicum* ATCC13032 using the cosmid pH79 (Hohn and Collins, Gene 11, 291-298 (1980)). O'Donohue (The Cloning and Molecular Analysis of Four Common Aromatic Amino Acid Biosynthetic Genes from *Corynebacterium glutamicum*. Ph.D. Thesis, National University of Ireland, Galway, 1997) describes the cloning of *C. glutamicum* genes using the  $\lambda$  Zap expression system described by Short et al. (Nucleic Acids Research, 16: 7583). To prepare a gene library of *C. glutamicum* in *E. coli* it is also possible to use plasmids such as pBR322 (Bolivar, Life Sciences, 25, 807-818 (1979)) or pUC9 (Vieira et al., 1982, Gene, 19:259-268). Suitable hosts are, in particular, those *E. coli* strains which are restriction- and recombination-defective. An example of these is the strain DH5 $\alpha$ mc $r$ , which has been described by Grant et al. (Proceedings of the National Academy of Sciences USA, 87 (1990) 4645-4649). The long DNA fragments cloned with the aid of cosmids can then in turn be subcloned and subsequently sequenced in the usual vectors

which are suitable for sequencing, such as is described e. g. by Sanger et al. (Proceedings of the National Academy of Sciences of the United States of America, 74:5463-5467, 1977).

- 5 The DNA sequences obtained can then be investigated with known algorithms or sequence analysis programs, such as e. g. that of Staden (Nucleic Acids Research 14, 217-232(1986)), the GCG program of Butler (Methods of Biochemical Analysis 39, 74-97 (1998)) the FASTA algorithm
- 10 of Pearson and Lipman (Proceedings of the National Academy of Sciences USA 85,2444-2448 (1988)) or the BLAST algorithm of Altschul et al. (Nature Genetics 6, 119-129 (1994)) and compared with the sequence entries which exist in databanks accessible to the public. Databanks for nucleotide
- 15 sequences which are accessible to the public are, for example, that of the European Molecular Biology Laboratories (EMBL, Heidelberg, Germany) or that of the National Center for Biotechnology Information (NCBI, Bethesda, MD, USA).
- 20 The invention provides a new DNA sequence of *C. glutamicum* which codes for the *opcA* gene and which is a constituent of the present invention as SEQ ID NO 1 and SEQ ID NO 4. The amino acid sequence of the corresponding protein has furthermore been derived from the present DNA sequence by
- 25 the methods described above. The resulting amino acid sequence of the *OpcA* gene product is shown in SEQ ID NO 3 and SEQ ID NO 5. The molecular weight resulting from the amino acid sequence of the *OpcA* gene product is approx. 34.7 kilo dalton.
- 30 SEQ ID NO 1 also shows the coding region of the *zwf* gene. The resulting amino acid sequence of the *Zwf* gene product is shown in SEQ ID NO 2. The molecular weight resulting from the amino acid sequence of the *Zwf* gene product is approx. 57.5 kilo dalton.

A gene library produced in the manner described above can furthermore be investigated by hybridization with nucleotide probes of known sequence, such as, for example, the zwf gene (JP-A-09224661). The cloned DNA of the clones  
5 which show a positive reaction in the hybridization is sequenced in turn to give on the one hand the known nucleotide sequence of the probe employed and on the other hand the adjacent new DNA sequences.

The invention also provides a new DNA sequence of C. glutamicum which codes for the opcA gene and which is a  
10 constituent of the present invention as SEQ ID NO 6 and SEQ ID NO 9. The amino acid sequence of the corresponding protein has furthermore been derived from the present DNA sequence by the methods described above. The resulting  
15 amino acid sequence of the OpcA gene product is shown in SEQ ID NO 8 and SEQ ID NO 10. The molecular weight resulting from the amino acid sequence of the OpcA gene product is approx. 34.7 kilo dalton.

SEQ ID NO 6 also shows the coding region of the zwf gene.  
20 The resulting amino acid sequence of the Zwf gene product is shown in SEQ ID NO 7. The molecular weight resulting from the amino acid sequence of the Zwf gene product is approx. 57.5 kilo dalton.

Another procedure for at least partly determining the amino  
25 acid sequence of the OpcA protein and the Zwf protein comprises purifying the glucose 6-phosphate dehydrogenase enzyme protein to homogeneity by chromatographic methods. Methods and instructions for protein purification and preparation are described e.g. in the textbook by Schleifer  
30 and Wensink: Practical Methods in Molecular Biology (Springer Verlag, Berlin, Germany, 1981), in the handbook by Harris and Ängal: Protein Purification Methods: A Practical Approach (IRL Press, Oxford, UK, 1989), in the  
textbook by Scopes: Protein Purification: Principles and  
35 Practice, 3<sup>rd</sup> ed. (Springer Verlag, New York, USA, 1993)

and in generally known textbooks and handbooks. The N-terminal amino acid sequence of the purified polypeptides can be determined by the method of N-terminal sequencing described by Edman (Archives of Biochemistry 22, 475 (1949)). Further methods and instructions for protein sequencing are described e. g. in Smith: Protein Sequencing Protocols: Methods in Molecular Biology, Vol. 64 and Vol. 112 (Humana Press, Totowa, NJ, USA, 1996) and in Kamp et al.: Protein Structure Analysis: Preparation, Characterization, and Microsequencing (Springer Verlag, New York, NY, USA, 1997).

It was possible to show in this manner that the enzyme glucose 6-phosphate dehydrogenase consists of two sub-units with in each case a molecular weight of approx. 30 kDa and approx. 60 kDa. The N-terminal amino acid sequence of the OpcA sub-unit and of the OpcA protein is shown in SEQ-ID-NO. 12. The N-terminal amino acid sequence of the Zwf sub-unit and of the Zwf protein is shown in SEQ-ID-NO. 11.

Coding DNA sequences which result from SEQ ID NO 1, SEQ ID NO 4, SEQ ID NO 6 or SEQ ID NO 9 by the degeneracy of the genetic code are also a constituent of the invention. In the same way, DNA sequences which hybridize with SEQ ID NO 4 or SEQ ID NO 9 or parts of SEQ ID NO 4 or SEQ ID NO 9 are a constituent of the invention. Conservative amino acid exchanges, such as e. g. exchange of glycine for alanine or of aspartic acid for glutamic acid in proteins, are furthermore known among experts as "sense mutations" which do not lead to a fundamental change in the activity of the protein, i.e. are of neutral function. It is furthermore known that changes on the N and/or C terminus of a protein cannot substantially impair or can even stabilize the function thereof. Information in this context can be found by the expert, inter alia, in Ben-Bassat et al. (Journal of Bacteriology 169:751-757 (1987)), in O'Regan et al. (Gene

77:237-251 (1989)), in Sahin-Toth et al. (Protein Sciences 3:240-247 (1994)), in Hochuli et al. (Bio/Technology 6:1321-1325 (1988)) and in known textbooks of genetics and molecular biology. Amino acid sequences which result in a  
5 corresponding manner from SEQ ID NO 3, SEQ ID NO 5, SEQ ID NO 8 or SEQ ID NO 10 are also a constituent of the invention.

Finally, DNA sequences which are prepared by the polymerase chain reaction (PCR) using primers which result from SEQ ID  
10 NO 4 or SEQ ID NO 9 are a constituent of the invention. Such oligonucleotides typically have a length of at least 15 nucleotides.

Instructions for identifying DNA sequences by means of hybridization can be found by the expert, inter alia, in  
15 the handbook "The DIG System Users Guide for Filter Hybridization" from Boehringer Mannheim GmbH (Mannheim, Germany, 1993) and in Liebl et al. (International Journal of Systematic Bacteriology (1991) 41: 255-260).

Instructions for amplification of DNA sequences with the  
20 aid of the polymerase chain reaction (PCR) can be found by the expert, inter alia, in the handbook by Gait: Oligonukleotide synthesis: a practical approach (IRL Press, Oxford, UK, 1984) and in Newton and Graham: PCR (Spektrum Akademischer Verlag, Heidelberg, Germany, 1994).

25 The inventors have found that coryneform bacteria produce amino acids, in particular L-lysine, in an improved manner after over-expression of the opcA gene, optionally together with the zwf gene.

To achieve an over-expression, the number of copies of the  
30 corresponding genes can be increased, or the promoter and regulation region or the ribosome binding site upstream of the structural gene can be mutated. Expression cassettes which are incorporated upstream of the structural gene act in the same way. By inducible promoters, it is additionally

possible to increase the expression in the course of fermentative L-lysine production. The expression is likewise improved by measures to prolong the life of the mRNA. Furthermore, the enzyme activity is also increased by preventing the degradation of the enzyme protein. The genes or gene constructs can either be present in plasmids with a varying number of copies, or can be integrated and amplified in the chromosome. Alternatively, an over-expression of the genes in question can furthermore be achieved by changing the composition of the media and the culture procedure.

Instructions in this context can be found by the expert, inter alia, in Martin et al. (Bio/Technology 5, 137-146 (1987)), in Guerrero et al. (Gene 138, 35-41 (1994)), Tsuchiya and Morinaga (Bio/Technology 6, 428-430 (1988)), in Eikmanns et al. (Gene 102, 93-98 (1991)), in European Patent Specification EPS 0 472 869, in US Patent 4,601,893, in Schwarzer and Pühler (Bio/Technology 9, 84-87 (1991)), in Reinscheid et al. (Applied and Environmental Microbiology 60, 126-132 (1994)), in LaBarre et al. (Journal of Bacteriology 175, 1001-1007 (1993)), in Patent Application WO 96/15246, in Japanese Laid-Open Specification JP-A-10-229891, in Jensen and Hammer (Biotechnology and Bioengineering 58, 191-195 (1998)), in Makrides (Microbiological Reviews 60:512-538 (1996)) and in known textbooks of genetics and molecular biology.

By way of example, the *opcA* gene according to the invention was over-expressed with the aid of plasmids.

Suitable plasmids are those which are replicated in coryneform bacteria. Numerous known plasmid vectors, such as e. g. pZ1 (Menkel et al., Applied and Environmental Microbiology (1989) 64: 549-554), pEKEX1 (Eikmanns et al., Gene 102:93-98 (1991)) or pHS2-1 (Sonnen et al., Gene 107:69-74 (1991)) are based on the cryptic plasmids

pHM1519, pBL1 or pGA1. Other plasmid vectors, such as e. g. those based on pCG4 (US-A 4,489,160), or pNG2 (Serwold-Davis et al., FEMS Microbiology Letters 66, 119-124 (1990)), or pAG1 (US-A 5,158,891), can be used in the same  
5 manner.

The E. coli - C. glutamicum shuttle vector pEC-T18mob2 shown in Figure 2 was used as an example. After incorporation of the opca gene and the zwf gene into the SphI/SalI cleavage site region of pEC-T18mob2, the plasmid  
10 pECzwfopca shown in Figure 3 was formed.

Plasmid vectors which are furthermore suitable are also those with the aid of which the process of gene amplification by integration into the chromosome can be used, as has been described, for example, by Reinscheid et al. (Applied and Environmental Microbiology 60, 126-132  
15 (1994)) for duplication or amplification of the hom-thrB operon. In this method, the complete gene is cloned in a plasmid vector which can replicate in a host (typically E. coli), but not in C. glutamicum. Possible vectors are, for example, pSUP301 (Simon et al., Bio/Technology 1, 784-791  
20 (1983)), pK18mob or pK19mob (Schäfer et al., Gene 145, 69-73 (1994)), pGEM-T (Promega corporation, Madison, WI, USA), pCR2.1-TOPO (Shuman (1994). Journal of Biological Chemistry 269:32678-84; US-A 5,487,993), pCR®Blunt  
25 (Invitrogen, Groningen, Holland; Bernard et al., Journal of Molecular Biology, 234: 534-541 (1993)), pEM1 (Schrumpf et al, 1991, Journal of Bacteriology 173:4510-4516) or pBGS8 (Spratt et al., 1986, Gene 41: 337-342). The plasmid vector which contains the gene to be amplified is then transferred  
30 into the desired strain of C. glutamicum by conjugation or transformation. The method of conjugation is described, for example, by Schäfer et al. (Applied and Environmental Microbiology 60, 756-759 (1994)). Methods for transformation are described, for example, by Thierbach et al. (Applied Microbiology and Biotechnology 29, 356-362  
35

(1988)), Dunican and Shivnan (Bio/Technology 7, 1067-1070 (1989)) and Tauch et al. (FEMS Microbiological Letters 123, 343-347 (1994)). After homologous recombination by means of a "cross over" event, the resulting strain contains at least two copies of the gene in question.

In addition, it may be advantageous for the production of amino acids, in particular L-lysine, to amplify or over-express one or more enzymes of the particular biosynthesis pathway, of glycolysis, of anaplerosis, of the pentose phosphate pathway or of amino acid export, in addition to the opcA gene, optionally together with the zwf gene.

Thus, for example, for the preparation of L-lysine, it may be advantageous for one or more genes chosen from the group consisting of

- the dapA gene which codes for dihydrodipicolinate synthase (EP-B 0 197 335),
- the lysC gene which codes for a feed back resistant aspartate kinase (Kalinowski et al. (1990), Molecular and General Genetics 224: 317-324),
- the gap gene which codes for glycerolaldehyde 3-phosphate dehydrogenase (Eikmanns (1992), Journal of Bacteriology 174:6076-6086),
- the pyc gene which codes for pyruvate carboxylase (DE-A-198 31 609),
- the tkt gene which codes for transketolase (accession number AB023377 of the databank of European Molecular Biology Laboratories (EMBL, Heidelberg, Germany)),
- the gnd gene which codes for 6-phosphogluconate dehydrogenase (JP-A-9-224662),
- the lysE gene which codes for lysine export (DE-A-195 48 222),



- the zwal gene (DE 199 59 328.0; DSM 13115), or
- the eno gene which codes for enolase (DE: 199 41 478.5),
- the tal gene which codes for transaldolase (DSM 13263)

to be amplified, in particular over-expressed, at the same  
5 time.

It may furthermore be advantageous for the production of amino acids, in particular L-lysine, at the same time to attenuate

- the pck gene which codes for phosphoenol pyruvate  
10 carboxykinase (DE 199 50 409.1 DSM 13047) and/or
- the pgi gene which codes for glucose 6-phosphate isomerase (US 09/396,478, DSM 12969), or
- the poxB gene which codes for pyruvate oxidase (DE 199 51 975.7; DSM 13114), or
- 15 • the zwa2 gene (DE: 199 59 327.2; DSM 13113)

in addition to the amplification of the opcA gene, optionally in combination with the zwf gene.

In addition to over-expression of the opcA gene it may furthermore be advantageous for the production of amino  
20 acids, in particular L-lysine, to eliminate undesirable side reactions (Nakayama: "Breeding of Amino Acid Producing Micro-organisms", in: Overproduction of Microbial Products, Krumphanzl, Sikyta, Vanek (eds.), Academic Press, London, UK, 1982).

25 The microorganisms prepared according to the invention can be cultured continuously or discontinuously in the batch process (batch culture) or in the fed batch (feed process) or repeated fed batch process (repetitive feed process) for the purpose of production of L-amino acids, in particular

L-lysine. A summary of known culture methods are described in the textbook by Chmiel (Bioprozesstechnik 1. Einführung in die Bioverfahrenstechnik [Bioprocess Technology 1. Introduction to Bioprocess Technology (Gustav Fischer Verlag, Stuttgart, 1991)) or in the textbook by Storhas (Bioreaktoren und periphere Einrichtungen [Bioreactors and Peripheral Equipment] (Vieweg Verlag, Braunschweig/Wiesbaden, 1994)).

The culture medium to be used must meet the requirements of the particular strains in a suitable manner. Descriptions of culture media for various microorganisms are contained in the handbook "Manual of Methods for General Bacteriology" of the American Society for Bacteriology (Washington D.C., USA, 1981). Sugars and carbohydrates, such as e. g. glucose, sucrose, lactose, fructose, maltose, molasses, starch and cellulose, oils and fats, such as e. g. soya oil, sunflower oil, groundnut oil and coconut fat, fatty acids, such as e. g. palmitic acid, stearic acid and linoleic acid, alcohols, such as e. g. glycerol and ethanol, and organic acids, such as e. g. acetic acid, can be used as the source of carbon. These substance can be used individually or as a mixture. Organic nitrogen-containing compounds, such as peptones, yeast extract, meat extract, malt extract, corn steep liquor, soya bean flour and urea, or inorganic compounds, such as ammonium sulphate, ammonium chloride, ammonium phosphate, ammonium carbonate and ammonium nitrate, can be used as the source of nitrogen. The sources of nitrogen can be used individually or as a mixture. Phosphoric acid, potassium dihydrogen phosphate or dipotassium hydrogen phosphate or the corresponding sodium-containing salts can be used as the source of phosphorus. The culture medium must furthermore comprise salts of metals, such as e. g. magnesium sulfate or iron sulfate, which are necessary for growth. Finally, essential growth substances, such as amino acids and vitamins, can be employed in addition to the

above-mentioned substances. Suitable precursors can moreover be added to the culture medium. The starting substances mentioned can be added to the culture in the form of a single batch, or can be fed in during the culture in a suitable manner.

Basic compounds, such as sodium hydroxide, potassium hydroxide, ammonia or aqueous ammonia, or acid compounds, such as phosphoric acid or sulfuric acid, can be employed in a suitable manner to control the pH. Antifoams, such as e. g. fatty acid polyglycol esters, can be employed to control the development of foam. Suitable substances having a selective action, such as e. g. antibiotics, can be added to the medium to maintain the stability of plasmids. To maintain aerobic conditions, oxygen or oxygen-containing gas mixtures, such as e. g. air, are introduced into the culture. The temperature of the culture is usually 20°C to 45°C, and preferably 25°C to 40°C. Culturing is continued until a maximum of L-amino acid has formed. This target is usually reached within 10 hours to 160 hours.

The analysis of L-amino acids can be carried out by anion exchange chromatography with subsequent ninhydrin derivatization, as described by Spackman et al. (Analytical Chemistry, 30, (1958), 1190).

The following microorganism has been deposited at the Deutsche Sammlung für Mikroorganismen und Zellkulturen (DSMZ = German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany) in accordance with the Budapest Treaty:

- *Corynebacterium glutamicum* ATCC13032/pECzwfopCA as DSM 13264

SEQ ID NO 1 also contains the new devB gene. The process according to the invention is used for the fermentative preparation of amino acids, in particular L-lysine.

Sequences attached:

The following sequences are attached in the form of a sequence protocol:

	SEQ ID NO:	Description:
5	1	DNA sequence isolated from <i>Corynebacterium glutamicum</i> ATCC13032
	2	Amino acid sequence of the Zwf protein derived from SEQ ID NO 1
	3	Amino acid sequence of the OpcA protein derived from SEQ ID NO 1
10	4	DNA sequence of the <i>opcA</i> gene of ATCC13032 taken from SEQ ID NO 1
	5	Amino acid sequence of the OpcA protein derived from SEQ ID NO 4
15	6	DNA sequence isolated from <i>Corynebacterium glutamicum</i> AS019
	7	Amino acid sequence of the Zwf protein derived from SEQ ID NO 6
	8	Amino acid sequence of the OpcA protein derived from SEQ ID NO 6
20	9	DNA sequence of the <i>opcA</i> gene of AS019 taken from SEQ ID NO 6
	10	Amino acid sequence of the OpcA protein derived from SEQ ID NO 9
25	11	Amino acid sequence of the N-terminus of the Zwf protein of the glucose 6-phosphate dehydrogenase from ATCC13032 which can be isolated
30	12	Amino acid sequence of the N-terminus of the OpcA protein of the glucose 6-phosphate dehydrogenase, which can be isolated, from ATCC13032

The following figures are attached:

Figure 1: Map of the plasmid pBOB102

Figure 2: Map of the plasmid pEC-T18mob2

Figure 3: Map of the plasmid pECzwfopcA

5 The abbreviations used have the following meaning:

Figure 1:

Neo r :	Neomycin/kanamycin resistance
ColE1 ori:	origin of replication of plasmid ColE1
CMV:	Cytomegalovirus promoter
10 lacP:	promotor of lac operon
lacZ:	5'-end of $\beta$ -galactosidase gene (lacZa gene fragment)
SV40 3' splice	3' splice site of Simian Virus 40
SV40 polyA:	polyadenylation site of Simian Virus 40
15 fl(-)ori:	origin of replication of filamentousphage fl
SV40 ori:	origin of replication of Simian Virus 40

Figures 2 and 3:

20 Tet:	Resistance gene for tetracycline
oriV:	Plasmid-coded replication origin of E. coli
RP4mob:	mob region for mobilizing the plasmid
rep:	Plasmid-coded replication origin from C. glutamicum plasmid pGA1
25 per:	Gene for controlling the number of copies from PGA1
lacZ-alpha:	lacZ $\alpha$ gene fragment (N-terminus) of the $\beta$ -galactosidase gene
lacZalpha':	5'-Terminus of the lacZ $\alpha$ gene fragment
30 'lacZalpha:	3'-Terminus of the lacZ $\alpha$ gene fragment
zwf:	zwf gene
opcA:	opcA gene

## Further abbreviations:

	ApaI:	cleavage site of restriction enzyme ApaI
	BamHI:	cleavage site of restriction enzyme BamHI
	ClaI:	cleavage site of restriction enzyme ClaI
5	EcoRI:	cleavage site of restriction enzyme EcoRI
	HindIII:	cleavage site of restriction enzyme HindIII
	MstII:	cleavage site of restriction enzyme MstII
	NheI:	cleavage site of restriction enzyme NheI
	NsiI:	cleavage site of restriction enzyme NsiI
10	SacI:	cleavage site of restriction enzyme SacI
	SalI:	cleavage site of restriction enzyme SalI
	SpeI:	cleavage site of restriction enzyme SpeI
	SphI:	cleavage site of restriction enzyme SphI
	SspI:	cleavage site of restriction enzyme SspI
15	XbaI:	cleavage site of restriction enzyme XbaI

### Examples

The following examples will further illustrate this invention. The molecular biology techniques, e.g. plasmid DNA isolation, restriction enzyme treatment, ligations, standard transformations of *Escherichia coli* etc. used are, (unless stated otherwise), described by Sambrook et al., (Molecular Cloning. A Laboratory Manual (1989) Cold Spring Harbour Laboratories, USA).

#### Example 1

- 10 Construction of a gene library of *Corynebacterium glutamicum* strain AS019

A DNA library of *Corynebacterium glutamicum* strain AS019 (Yoshihama et al., Journal of Bacteriology 162, 591-597 (1985)) was constructed using  $\lambda$  Zap Express™ system, (Short et al., (1988) Nucleic Acids Research, 16: 7583-7600), as described by O'Donohue (O'Donohue, M. (1997). The Cloning and Molecular Analysis of Four Common Aromatic Amino Acid Biosynthetic Genes from *Corynebacterium glutamicum*. Ph.D. Thesis, National University of Ireland, Galway.).  $\lambda$  Zap Express™ kit was purchased from Stratagene (Stratagene, 11011 North Torrey Pines Rd., La Jolla, California 92037.) and used according to the manufacturers instructions. AS019-DNA was digested with restriction enzyme Sau3A and ligated to BamHI treated and dephosphorylated  $\lambda$  Zap Express™ arms.

#### Example 2

Cloning and sequencing of the *opcA* and *zwf* gene

##### 1. Construction of a *zwf* probe

A radiolabelled oligonucleotide, internal to the *zwf* gene, was used to probe the AS019  $\lambda$  Zap Express™ library described above. The oligonucleotide was produced using degenerate PCR primers internal to the *zwf* gene. The

degenerate nucleotide primers designed for the PCR amplification of zwf DNA fragments were as follows:

zwf1: 5' ATY GAY CAC TAY YTS GGY AAR GA 3'

zwf2: 5' RAA WGG MAC RCC YKS CCA 3'

5 with R=A+G; Y=C+T; W=A+T; M=A+C; S=G+C; K=T+G.

The estimated size of the resulting PCR product was 480bp approximately.

Optimal PCR conditions were determined to be as follows:

- 35 cycles
  - 10 94°C for 1 minute
  - 60°C for 1 minute
  - 72°C for 30 seconds
  - 2.5 - 3.5 mM MgCl<sub>2</sub>
  - 100 - 150 ng AS019 genomic DNA
- 15 Sequence analysis of the resulting PCR product confirmed the product to be an internal portion of the zwf gene. Sequence analysis was carried out using the universal forward and reverse primers, and T7 sequencing kit from Pharmacia Biotech, (St. Albans, Herts, UK).
- 20 2. Cloning
- Screening of the AS019  $\lambda$  Zap Express<sup>TM</sup> library was carried out according to the  $\lambda$  Zap Express<sup>TM</sup> system protocol, (Stratagene, 11011 North Torrey Pines Rd., La Jolla, California 92037.). Southern Blot analysis was then carried
- 25 out on isolated clones. Southern transfer of DNA was as described in the Schleicher and Schuell protocols manual employing Nytran<sup>TM</sup> as membrane („Nytran, Modified Nylon-66 Membrane Filters“ (March 1987), Schleicher and Schuell, Dassel, Germany). Double stranded DNA fragments, generated
- 30 using the same primers and optimal PCR conditions as described above, were radiolabelled with  $\alpha$ -<sup>32</sup>P-dCTP using the Multiprime<sup>TM</sup> DNA labelling kit from Amersham Life



Science (Amersham Pharmacia Biotech UK Limited, Little Chalfont, Buckinghamshire, UK) according to the manufacturers instructions. Prehybridisation, hybridization and washing conditions were as described in the Schleicher and Schuell protocols manual. Autoradiography was carried out according to the procedure outlined in the handbook of Sambrook et al. using AgFa Curix RPIL film. Thus several zwf clones were identified. Plasmid DNA was isolated from one of the clones, designated pBOB102 (Figure 1) and chosen for further analysis.

### 3. Sequencing

The Sanger Dideoxy chain termination method of Sanger et al. (Proceedings of the National Academy of Sciences USA 74, 5463-5467 (1977)) was used to sequence the cloned insert of pBOB102. The method was applied using the T7 sequencing kit and  $\alpha$ -<sup>35</sup>S-dCTP from Pharmacia Biotech (St. Albans, Herts, UK). Samples were electrophoresed for 3-8 hours on 6% polyacrylamide/urea gels in TBE buffer at a constant current of 50 mA, according to the Pharmacia cloning and sequencing instructions manual („T7 Sequencing™ Kit“, ref.XY-010-00-19, Pharmacia Biotech, 1994). Initial sequence analysis was carried out using the universal forward and M13 reverse primers obtained from Pharmacia Biotech:

#### Universal forward

primer: 5' GTA ATA CGA CTC ACT ATA GGG C 3'

M13 reverse primer: 5' GGA AAC AGC TAT GAC CAT G 3'

Internal primers were subsequently designed from the sequence obtained which allowed the entire opcA gene to be deduced. The sequences of the internal primers were as follows:

Internal primer 1: 5' TCA ACC CTG AGT CCA CC 3'

Internal primer 2: 5' CTG ACC ACG AGC GGA GG 3'

Internal primer 3: 5' ATG GTG ATC TGG ACG TG 3'

Internal primer 4: 5' CTG GCG ACT TGG CTC GA 3'  
Internal primer 5: 5' CTT CCG GAT ACC ACC ACC 3'

Sequence obtained was then analyzed using the DNA Strider programme, (Marck (1988), Nucleic Acids Research 16: 1829-1836), version 1.0 on an Apple Macintosh computer. This program allowed for analyses such as restriction site usage, open reading frame analysis and codon usage determination. Searches between DNA sequence obtained and those in EMBL and Genbank databases were achieved using the BLAST programme, (Altschul et al., (1997) Nucleic Acids Research, 25: 3389-3402). DNA and protein sequences were aligned using the Clustal V and Clustal W programs (Higgins and Sharp, 1988 Gene 73: 237-244).

The sequence thus obtained is shown in SEQ ID NO 6. The analysis of the nucleotide sequence obtained revealed an open reading frame of 957 base pairs which was designated as opcA gene. It codes for a protein of 319 amino acids shown in SEQ ID NO 8 and SEQ ID NO 10. The coding region of the zwf gene is also shown in SEQ ID NO 6. The amino acid sequence of the Zwf-Protein composed of 514 amino acids is shown in SEQ ID NO 7.

### Example 3

Preparation of a genomic cosmid gene library from *Corynebacterium glutamicum* ATCC 13032

Chromosomal DNA from *Corynebacterium glutamicum* ATCC 13032 was isolated as described by Tauch et al. (1995, Plasmid 33:168-179) and partly cleaved with the restriction enzyme Sau3AI (Amersham Pharmacia, Freiburg, Germany, Product Description Sau3AI, Code no. 27-0913-02). The DNA fragments were dephosphorylated with shrimp alkaline phosphatase (Roche Molecular Biochemicals, Mannheim, Germany, Product Description SAP, Code no. 1758250). The DNA of the cosmid vector SuperCos1 (Wahl et al. (1987) Proceedings of the

National Academy of Sciences USA 84:2160-2164), obtained from Stratagene (La Jolla, USA, Product Description SuperCos1 Cosmid Vector Kit, Code no. 251301) was cleaved with the restriction enzyme XbaI (Amersham Pharmacia, Freiburg, Germany, Product Description XbaI, Code no. 27-0948-02) and likewise dephosphorylated with shrimp alkaline phosphatase. The cosmid DNA was then cleaved with the restriction enzyme BamHI (Amersham Pharmacia, Freiburg, Germany, Product Description BamHI, Code no. 27-0868-04).

10 The cosmid DNA treated in this manner was mixed with the treated ATCC13032 DNA and the batch was treated with T4 DNA ligase (Amersham Pharmacia, Freiburg, Germany, Product Description T4-DNA-Ligase, Code no. 27-0870-04). The ligation mixture was then packed in phages with the aid of

15 Gigapack II XL Packing Extracts (Stratagene, La Jolla, USA, Product Description Gigapack II XL Packing Extract, Code no. 200217). For infection of the E. coli strain NM554 (Raleigh et al. 1988, Nucleic Acid Research 16:1563-1575) the cells were taken up in 10 mM MgSO<sub>4</sub> and mixed with an

20 aliquot of the phage suspension. The infection and titering of the cosmid library were carried out as described by Sambrook et al. (1989, Molecular Cloning: A laboratory Manual, Cold Spring Harbor), the cells being plated out on LB agar (Lennox, 1955, Virology, 1:190) with 100 µg/ml

25 ampicillin. After incubation overnight at 37°C, recombinant individual clones were selected.

#### Example 4

Isolation and sequencing of the opca and zwf gene of ATCC 13032

30 The cosmid DNA of an individual colony was isolated with the Qiaprep Spin Miniprep Kit (Product No. 27106, Qiagen, Hilden, Germany) in accordance with the manufacturer's instructions and partly cleaved with the restriction enzyme Sau3AI (Amersham Pharmacia, Freiburg, Germany, Product

35 Description Sau3AI, Product No. 27-0913-02). The DNA

fragments were dephosphorylated with shrimp alkaline phosphatase (Roche Molecular Biochemicals, Mannheim, Germany, Product Description SAP, Product No. 1758250). After separation by gel electrophoresis, the cosmid

5 fragments in the size range of 1500 to 2000 bp were isolated with the QiaExII Gel Extraction Kit (Product No. 20021, Qiagen, Hilden, Germany). The DNA of the sequencing vector pZero-1, obtained from Invitrogen (Groningen, Holland, Product Description Zero Background Cloning Kit,

10 Product No. K2500-01) was cleaved with the restriction enzyme BamHI (Amersham Pharmacia, Freiburg, Germany, Product Description BamHI, Product No. 27-0868-04). The ligation of the cosmid fragments in the sequencing vector pZero-1 was carried out as described by Sambrook et al.

15 (1989, Molecular Cloning: A laboratory Manual, Cold Spring Harbor), the DNA mixture being incubated overnight with T4 ligase (Pharmacia Biotech, Freiburg, Germany). This ligation mixture was then electroporated (Tauch et al. 1994, FEMS Microbiol. Letters, 123:343-7) into the E. coli strain DH5 $\alpha$ MCR (Grant, 1990, Proceedings of the National

20 Academy of Sciences U.S.A., 87:4645-4649) and plated out on LB agar (Lennox, 1955, Virology, 1:190) with 50  $\mu$ g/ml zeocin. The plasmid preparation of the recombinant clones was carried out with Biorobot 9600 (Product No. 900200,

25 Qiagen, Hilden, Germany). The sequencing was carried out by the dideoxy chain-stopping method of Sanger et al. (1977, Proceedings of the National Academy of Sciences U.S.A., 74:5463-5467) with modifications according to Zimmermann et al. (1990, Nucleic Acids Research, 18:1067). The "RR

30 dRhodamin Terminator Cycle Sequencing Kit" from PE Applied Biosystems (Product No. 403044, Weiterstadt, Germany) was used. The separation by gel electrophoresis and analysis of the sequencing reaction were carried out in a "Rotiphoresis NF Acrylamide/Bisacrylamide" Gel (29:1) (Product No.

35 Al24.1, Roth, Karlsruhe, Germany) with the "ABI Prism 377" sequencer from PE Applied Biosystems (Weiterstadt, Germany).

The raw sequence data obtained were then processed using the Staden program package (1986, Nucleic Acids Research, 14:217-231) version 97-0. The individual sequences of the pZerol derivatives were assembled to a continuous contig.

5 The computer-assisted coding region analysis were prepared with the XNIP program (Staden, 1986, Nucleic Acids Research, 14:217-231). Further analyses were carried out with the "BLAST search programs" (Altschul et al., 1997, Nucleic Acids Research, 25:3389-3402), against the non-

10 redundant databank of the "National Center for Biotechnology Information" (NCBI, Bethesda, MD, USA).

The nucleotide sequence obtained is shown in SEQ ID NO 1. Analysis of the nucleotide sequence showed a coding region of 957 base pairs, which was called the opcA gene. The opcA

15 gene, including its stop codon, is shown in SEQ ID NO 4. The opcA gene codes for a protein of 319 amino acids shown in SEQ ID NO 3 and SEQ ID NO 5.

#### Example 5

Purification and N-terminal sequencing of the glucose-6-phosphate dehydrogenase of *Corynebacterium glutamicum*

20 ATCC13032.

##### 1. Culture of strain ATCC 13032

For purification of the glucose-6-phosphate dehydrogenase *Corynebacterium glutamicum* ATCC 13032 was grown aerobically

25 on minimal medium at 30°C in a Labfors fermentation system (Infors AG, Bottmingen, Switzerland). A preculture (Bacto® Brain Heart Infusion medium, Difco Laboratories, Detroit, USA) was incubated for 15 hours at 30°C and used for inoculation of 2.5 l minimal medium. The medium contained

30 the following constituents (amounts per liter): 20 g  $(\text{NH}_4)_2\text{SO}_4$ ; 1 g  $\text{KH}_2\text{PO}_4$ ; 1 g  $\text{K}_2\text{HPO}_4$ ; 0.25 g  $\text{MgSO}_4 \times 7 \text{ H}_2\text{O}$ ; 10 mg  $\text{CaCl}_2$ ; 0.2 mg biotin; 30 mg protocatechuic acid; 1 mg  $\text{FeSO}_4 \times 7 \text{ H}_2\text{O}$ ; 1 mg  $\text{MnSO}_4 \times \text{H}_2\text{O}$ ; 0.1 mg  $\text{ZnSO}_4 \times 7 \text{ H}_2\text{O}$ ; 0.02 mg  $\text{CuSO}_4$ ; 0.002 mg  $\text{NiCl}_2 \times 6 \text{ H}_2\text{O}$ ; 1.2 g  $\text{HCl}$ ; 0.2 g

polypropylene glycol; 75 mg tritriplex II and 100 g glucose. During fermentation sodium hydroxide was continuously added in order to keep the pH-value constant at 7.0. The cells were harvested in the late exponential  
5 growth phase. After centrifugation using an Avanti J-25 centrifuge and a JA10 rotor of Beckman (Fullerton, USA) at 6400 g for 15 minutes at 4°C and washing in 100 mM TRIS-HCl pH 7.5 containing 10 mM MgCl<sub>2</sub> the sediment was stored at -20°C until use.

## 10 2. Enzyme purification

Disruption of cells was carried out in a disintegration system (Disintegrator S, BIOmatic, Rodgau-Hainhausen, Germany). The cells were previously resuspended in a pH 7.5 buffer consisting of 100 mM TRIS-HCl, 10 mM MgCl<sub>2</sub>, 0.75 mM  
15 DTT and a mixture of several protease inhibitors (complete™, Roche, Mannheim, Germany). The ratio of the cell wet weight to the total suspension weight was adjusted to 0.3. After addition of 100 ml glass beads with a diameter of 0.1 to 0.25 mm (Fisher scientific, Düsseldorf,  
20 Germany) per 100 ml total suspension volume, cell disruption was performed at 5000 rpm for 12 Minutes. A temperature increase during disruption was prevented by ice cooling. After removal of glass beads an ultracentrifugation step was carried through using an L8-70 M centrifuge and a  
25 Ti45 rotor of Beckman (Fullerton, USA) at 235000 g for 90 minutes at 4°C. The supernatant was used as crude extract for the purification of the glucose-6-phosphate dehydrogenase. All purification steps were carried out with a Biosys2000 system of Beckman (Fullerton, USA).

30 The crude extract was applied to an XK 50/30 column (Pharmacia, Freiburg, Germany), which contained Fractogel EMD DEAE-650(S) material (Merck, Darmstadt). The total bed volume was 500 ml. The column was previously equilibrated with 50 mM TRIS-HCl pH 7.5 containing 30 mM MgCl<sub>2</sub> and 0.75  
35 mM DTT. After application of the crude extract the column

was washed with the same buffer containing 144 mM KCl. Elution was performed within 95 minutes by a linear KCl gradient from 144 mM up to 320 mM. The flow rate was 7.4 ml/min. The active fractions were pooled and concentrated  
5 in centrprep® 30 concentrators (Amicon, Beverly, USA) using a Varifuge 3.0R centrifuge (Heraeus, Hanau, Germany) at 1500 g and 4°C. By dilution with 50 mM TRIS-HCl pH 7.5 containing 30 mM MgCl<sub>2</sub> and 0.75 mM DTT the KCl  
concentration was adjusted to 40 mM. After that the  
10 partially purified glucose-6-phosphate dehydrogenase was applied to an XK26/20 column (Pharmacia, Freiburg, Germany), which was filled with 65 ml Red-Sepharose CL6B (Pharmacia, Freiburg, Germany). The column was equilibrated with 50 mM TRIS-HCl pH 7.5 containing 30 mM MgCl<sub>2</sub> and 0.75  
15 mM DTT. Elution was carried out within 590 minutes by a linear 0 - 800 mM KCl gradient at a flow rate of 0.87 ml/min.

After pooling of the active glucose-6-phosphate dehydrogenase fractions, the KCl concentration was reduced  
20 to 10 mM in the same way as described above. After that the solution was applied to an XK16/20 column (Pharmacia, Freiburg, Germany), which contained 20 ml of a 2'5'-ADP-sepharose matrix (Pharmacia, Freiburg, Germany). The column was equilibrated with the same buffer as the Red-Sepharose  
25 CL6B column. Elution was performed by an 0 to 2 mM NADP linear gradient. The active glucose-6-phosphate dehydrogenase-fractions were pooled and applied to a gel filtration column.

For gel filtration a Superdex G200pg column (Pharmacia,  
30 Freiburg, Germany) with a diameter of 1,6 cm and a bed volume of 114 ml was used. The elution at a flow rate of 1 ml/min was carried through with 50 mM TRIS-HCl pH 7.5 containing 30 mM MgCl<sub>2</sub>, 200 mM KCl and 0.75 mM DTT. The active fractions were pooled and concentrated by  
35 ultrafiltration in centrprep® 30 concentrators (Amicon,

Beverly, USA). After addition of 50 % (v/v) glycerol to the purified glucose-6-phosphate dehydrogenase solution it was stored at -20°C.

During the whole purification process the glucose-6-phosphate-dehydrogenase activity and the protein concentration were measured.

The assay system for determination of the glucose-6-phosphate-dehydrogenase-activity contained 50 mM TRIS-HCl pH 7.5, 10 mM MgCl<sub>2</sub>, 1 mM NADP and 200 mM potassium glutamate. The reaction was initiated by addition of 4 mM glucose-6-phosphate and the formation of NADPH was followed by measuring the increase in absorbance at 340 nm at 30 °C. Protein concentrations were determined spectrophotometrically after Coomassie Brilliant Blue staining (Stoscheck, Methods in Enzymology 182, 50-68 (1990)). As protein standard bovine serum albumin was used. All measurements were carried out using a UV-160 A photometer (Shimadzu, Kyoto, Japan).

The purity of the glucose-6-phosphate dehydrogenase was tested by denaturing discontinuous SDS-gel electrophoresis according to the method of Laemmli (Laemmli, U.K., Nature 227, 680-685 (1970)). After the third purification step using 2'5'-ADP sepharose ligand affinity material two different proteins with molecular weights of ca. 60 kDa and 30 kDa could be obtained. These two proteins could not be separated by gel filtration chromatography. The specific activity of this preparation was determined as 213 U/mg protein.

### 3. N-terminal sequencing

N-terminal sequencing of the purified glucose-6-P dehydrogenase was performed according to the procedure of Edman (Edman and Begg, European Journal of Biochemistry 1, 80-91 (1967)) using a Procise® Protein Sequencing System (Applied Biosystems, Foster City, USA).



For the 60 kDa protein the following N-terminal sequence was obtained: Xaa Xaa Xaa Xaa Xaa Pro Xaa Xaa Trp Xaa Asn Pro Leu Arg Asp. It is also shown in SEQ ID No 11.

For the 30 kDa protein the following N-terminal sequence was obtained: Met Ile Phe Xaa Leu Pro Asp Xaa Xaa Xaa Gln Gln Ile Ser Lys. It is also shown in SEQ ID No 12.

#### Example 6

Cloning of the zwf and opcA genes into the pGEM T-vector.

PCR was used to amplify DNA fragments containing the entire zwf and opcA genes of *C. glutamicum* ATCC13032 and flanking upstream and downstream regions. PCR reactions were carried out using oligonucleotide primers designed from SEQ ID NO 1 and SEQ ID NO 6. Genomic DNA was isolated from *Corynebacterium glutamicum* ATCC13032 according to Heery and Dunican (Applied and Environmental Microbiology, 59: 791-799 (1993)) and used as template. The primers used were:

zwf fwd. primer: 5' AGA ATC AGC ACG CTG CAT CAG 3'

opcA rev. primer: 5' AGT ATG GTG CGC GTA CTA 3'

PCR parameters were as follows:

35 cycles  
95°C for 3 minutes  
94°C for 1 minute  
47°C for 1 minute  
72°C for 45 seconds  
2.5 mM MgCl<sub>2</sub>  
approx. 150-200 ng DNA template.

The PCR product obtained was cloned into the commercially available pGEM-T vector purchased from Promega Corp. (pGEM-T Easy Vector System 1, cat. no. A1360, Promega UK, Southampton) using *E. coli* strain JM109 (Yanisch-Perron et al., Gene 33: 103-119 (1985)) as a host.

Example 7

## Preparation of the shuttle vector pEC-T18mob2

The *E. coli* - *C. glutamicum* shuttle vector pEC-T18mob2 was constructed according to the prior art.

- 5 The vector contains the replication region rep of the plasmid pGA1 including the replication effector per (US-A- 5,175,108; Nesvera et al., Journal of Bacteriology 179, 1525-1532 (1997)), the tetracycline resistance-imparting tetA(2) gene of the plasmid pAG1 (US-A- 5,158,891; gene  
10 library entry at the National Center for Biotechnology Information (NCBI, Bethesda, MD, USA) with accession number AF121000), the replication region oriV of the plasmid pMB1 (Sutcliffe, Cold Spring Harbor Symposium on Quantitative Biology 43, 77-90 (1979)), the lacZ $\alpha$  gene fragment  
15 including the lac promoter and a multiple cloning site (mcs) (Norranders et al. Gene 26, 101-106 (1983)) and the mob region of the plasmid RP4 (Simon et al., (1983) Bio/Technology 1:784-791).

- The vector constructed was transformed in the *E. coli*  
20 strain DH5 $\alpha$  (Hanahan, In: DNA cloning. A practical approach. Vol. I. IRL-Press, Oxford, Washington DC, USA). Selection for plasmid-carrying cells was made by plating out the transformation batch on LB agar (Sambrook et al., Molecular cloning: a laboratory manual. 2<sup>nd</sup> Ed. Cold Spring  
25 Harbor Laboratory Press, Cold Spring Harbor, N.Y.), which had been supplemented with 5 mg/l tetracycline. Plasmid DNA was isolated from a transformant with the aid of the QIAprep Spin Miniprep Kit from Qiagen and checked by restriction with the restriction enzyme EcoRI and HindIII  
30 subsequent agarose gel electrophoresis (0.8%).

The plasmid was called pEC-T18mob2 and is shown in Figure 2. It is deposited in the form of the strain *Escherichia coli* K-12 strain DH5 $\alpha$ /pEC-T18mob2 at the

Deutsche Sammlung für Mikroorganismen und Zellkulturen  
(DSMZ = German Collection of Microorganisms and Cell  
Cultures, Braunschweig, Germany) as DSM 13244.

#### Example 8

- 5 Expression of glucose-6-phosphate dehydrogenase in  
Corynebacterium glutamicum

The entire zwf and opcA genes were subsequently isolated  
from the pGEM T-vector containing these genes (see Example  
6) on an SphI/SalI fragment and cloned into the lacZα  
10 SphI/SalI sites of the E. coli - C. glutamicum shuttle  
vector pEC-T18mob2 (see Example 7 and Figure 2). This  
shuttle vector contains two SphI sites. The first is  
situated within the multiple cloning site of lacZα and the  
second is situated within the gene conferring tetracycline  
15 resistance. Tetracycline (Sigma-Aldrich, PO Box 2424,  
Wimborne, Dorset BH21 7YR, UK) (5mg/l) was used therefore  
as a selective pressure as only those clones containing the  
intact tetracycline resistance gene would grow. This new  
construct was designated pECzwfopca (Figure 3). Restriction  
20 enzyme analysis with SacI (Boehringer Mannheim GmbH,  
Germany) revealed the correct orientation of the zwf and  
opcA genes in the lacZα gene of pEC-T18mob2 i. e.  
downstream the lac promotor. Corynebacterium glutamicum  
ATCC13032 (American Type Culture Collection, Manassas, VA,  
25 USA) was transformed with this construct and  
electrotransformants were selected on Luria agar  
supplemented with isopropyl-thiogalactopyranoside (IPTG),  
5-bromo-4-chloro-3-indolyl-galactopyranoside (XGAL) and  
tetracycline at concentrations of 1 mM, 0.02% and 5 mg/l  
30 respectively. Agar plates were incubated for 48 hours at  
30°C. Rapid plasmid preparations were carried out as  
described by O'Gara and Dunican, (Applied and  
Environmental Microbiology 61: 4477-4479 (1995)), and Sac I  
restriction confirmed the presence of required clones. One  
35 of the clones was designated ATCC13032/pECzwfopca.

Example 9

Preparation of amino acid producers with an amplified opcA gene

The L-lysine-producing strain *Corynebacterium glutamicum* DSM5715 is described in EP-B-0435132 and the L-threonine-producing strain *Brevibacterium flavum* DSM5399 is described in EP-B-0385940. Both strains are deposited at the Deutsche Sammlung für Mikroorganismen und Zellkulturen [German Collection of Microorganisms and Cell Cultures] in Braunschweig (Germany) in accordance with the Budapest Treaty.

The strains DSM5715 and DSM5399 were transformed with the plasmid pECzwfopca (Example 8) using the electroporation method described by Liebl et al., (FEMS Microbiology Letters, 53:299-303 (1989)) Selection of the transformants took place on LBHIS agar comprising 18.5 g/l brain-heart infusion broth, 0.5 M sorbitol, 5 g/l Bacto-tryptone, 2.5 g/l Bacto-yeast extract, 5 g/l NaCl and 18 g/l Bacto-agar, which had been supplemented with 5 mg/l tetracycline. Incubation was carried out for 2 days at 33°C.

The strains obtained in this way were called DSM5715/pECzwfopca and DSM5399/pECzwfopca.

Example 10

Preparation of L-threonine

The *C. glutamicum* strain DSM5399/pECzwfopca obtained in Example 9 was cultured in a nutrient medium suitable for the production of threonine and the threonine content in the culture supernatant was determined.

For this, the strain was first incubated on an agar plate with the corresponding antibiotic (brain-heart agar with tetracycline (5 mg/l)) for 24 hours at 33°C. Starting from

this agar plate culture, a preculture was seeded (10 ml medium in a 100 ml conical flask). The complete medium CgIII was used as the medium for the preculture.

Medium Cg III:

NaCl 2.5 g/l

Bacto-Peptone 10 g/l

Bacto-Yeast extract 10 g/l

Glucose (autoclaved separately) 2% (w/v)

The pH was brought to pH 7.4.

5 Tetracycline (5 mg/l) was added to this. The preculture was incubated for 16 hours at 33°C at 240 rpm on a shaking machine. A main culture was seeded from this preculture such that the initial OD (660nm) of the main culture was 0.1. Medium MM was used for the main culture.

Medium MM:

CSL (corn steep liquor) 5 g/l

MOPS (morpholinopropanesulfonic acid) 20 g/l

Glucose (autoclaved separately) 50 g/l

$(\text{NH}_4)_2\text{SO}_4$  25 g/l

$\text{KH}_2\text{PO}_4$  0.1 g/l

$\text{MgSO}_4 \cdot 7 \text{ H}_2\text{O}$  1.0 g/l

$\text{CaCl}_2 \cdot 2 \text{ H}_2\text{O}$  10 mg/l

$\text{FeSO}_4 \cdot 7 \text{ H}_2\text{O}$  10 mg/l

MnSO <sub>4</sub> * H <sub>2</sub> O	5.0mg/l
Biotin (sterile-filtered)	0.3 mg/l
Thiamine * HCl (sterile-filtered)	0.2 mg/l
L-Leucine (sterile-filtered)	0.1 g/l
CaCO <sub>3</sub>	25 g/l

The CSL, MOPS and the salt solution were brought to pH 7 with aqueous ammonia and autoclaved. The sterile substrate and vitamin solutions were then added, as well as the CaCO<sub>3</sub> autoclaved in the dry state.

- 5    Culturing was carried out in a 10 ml volume in a 100 ml conical flask with baffles. Tetracycline (5 mg/l) was added. Culturing was carried out at 33°C and 80% atmospheric humidity.

- 10    After 72 hours, the OD was determined at a measurement wavelength of 660 nm with a Biomek 1000 (Beckmann Instruments GmbH, Munich). The amount of threonine formed was determined with an amino acid analyzer from Eppendorf-BioTronik (Hamburg, Germany) by ion exchange chromatography and post-column derivatization with ninhydrin detection.

- 15    The result of the experiment is shown in Table 1.

Table 1

Strain	OD	L-Threonine g/l
DSM5399	12.3	0.74
DSM5399/pECzwfopcA	9.9	1.0

Example 11

## Preparation of L-lysine

The *C. glutamicum* strain DSM5715/pECzwfopca obtained in Example 9 was cultured in a nutrient medium suitable for the production of lysine and the lysine content in the culture supernatant was determined.

For this, the strain was first incubated on an agar plate with the corresponding antibiotic (brain-heart agar with tetracycline (5 mg/l)) for 24 hours at 33°C. Starting from this agar plate culture, a preculture was seeded (10 ml medium in a 100 ml conical flask). The complete medium CgIII was used as the medium for the preculture.

Medium Cg III:

NaCl 2.5 g/l

Bacto-Peptone 10 g/l

Bacto-Yeast extract 10 g/l

Glucose (autoclaved separately) 2% (w/v)

The pH was brought to pH 7.4.

Tetracycline (5 mg/l) was added to this. The preculture was incubated for 16 hours at 33°C at 240 rpm on a shaking machine. A main culture was seeded from this preculture such that the initial OD (660nm) of the main culture was 0.1. Medium MM was used for the main culture.

## Medium MM:

CSL (corn steep liquor)	5 g/l
MOPS (morpholinopropanesulfonic acid)	20 g/l
Glucose (autoclaved separately)	58 g/l
 (NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub>	 25 g/l
KH <sub>2</sub> PO <sub>4</sub>	0.1 g/l
MgSO <sub>4</sub> * 7 H <sub>2</sub> O	1.0 g/l
CaCl <sub>2</sub> * 2 H <sub>2</sub> O	10 mg/l
FeSO <sub>4</sub> * 7 H <sub>2</sub> O	10 mg/l
MnSO <sub>4</sub> * H <sub>2</sub> O	5.0mg/l
Biotin (sterile-filtered)	0.3 mg/l
Thiamine * HCl (sterile-filtered)	0.2 mg/l
L-Leucine (sterile-filtered)	0.1 g/l
CaCO <sub>3</sub>	25 g/l

The CSL, MOPS and the salt solution were brought to pH 7 with aqueous ammonia and autoclaved. The sterile substrate and vitamin solutions were then added, as well as the CaCO<sub>3</sub>,  
5 autoclaved in the dry state.

Culturing was carried out in a 10 ml volume in a 100 ml conical flask with baffles. Tetracycline (5 mg/l) was added. Culturing was carried out at 33°C and 80% atmospheric humidity.



After 72 hours, the OD was determined at a measurement wavelength of 660 nm with a Biomek 1000 (Beckmann Instruments GmbH, München). The amount of lysine formed was determined with an amino acid analyzer from Eppendorf-  
5 BioTronik (Hamburg, Germany) by ion exchange chromatography and post-column derivatization with ninhydrin detection.

The result of the experiment is shown in Table 2.

Table 2

Strain	OD	L-Lysine HCl g/l
DSM5715	10.8	16.0
DSM5715/pECzwfopcA	8.1	17.1


PCT

Original (for SUBMISSION) - printed on 03.07.2000 02:19:24 PM

990213 BT

0-1	Form - PCT/RO/134 (EASY) Indications Relating to Deposited Microorganism(s) or Other Biological Material (PCT Rule 13bis)	
0-1-1	Prepared using	PCT-EASY Version 2.90 (updated 08.03.2000)
0-2	International Application No.	
0-3	Applicant's or agent's file reference	990213 BT
1	The indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:	
1-1	page	18
1-2	line	24-30
1-3	Identification of Deposit	
1-3-1	Name of depositary institution	DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH
1-3-2	Address of depositary institution	Mascheroder Weg 1b, D-38124 Braunschweig, Germany
1-3-3	Date of deposit	26 January 2000 (26.01.2000)
1-3-4	Accession Number	DSMZ 13264
1-4	Additional Indications	NONE
1-5	Designated States for Which Indications are Made	all designated States
1-6	Separate Furnishing of Indications These indications will be submitted to the International Bureau later	NONE

## FOR RECEIVING OFFICE USE ONLY

0-4	This form was received with the international application: (yes or no)	yes
0-4-1	Authorized officer  C.A.J.A. PASCHE	

## FOR INTERNATIONAL BUREAU USE ONLY

0-5	This form was received by the international Bureau on:	
0-5-1	Authorized officer	

## Patent claims

1. An isolated polynucleotide from coryneform bacteria,  
comprising at least one polynucleotide sequence chosen  
5 from the group consisting of
  - a) polynucleotide which is identical to the extent of  
at least 70 % to a polynucleotide which codes for  
polypeptides which comprise at least one of the  
amino acid sequences according to SEQ ID No. 3 or  
10 SEQ ID No. 5 or SEQ ID No. 8 or SEQ ID No. 10,
  - b) polynucleotide which codes for polypeptides which  
comprise amino acid sequences which are identical  
to the extent of at least 70 % to the amino acid  
sequences according to SEQ ID No.3 or SEQ ID No. 5  
15 or according to SEQ ID No. 8 or SEQ ID No. 10,
  - c) polynucleotide which is complementary to the  
polynucleotides of a) or b), or
  - d) polynucleotide comprising at least 15 successive  
nucleotides of the polynucleotide sequences of a),  
20 b) or c).
2. A polynucleotide as claimed in claim 1  
wherein the polynucleotide is a preferably recombinant  
DNA which is capable of replication in coryneform  
bacteria and additionally contains at least one of the  
25 nucleotide sequences which code for the genes tal,  
tkt, zwf and devB.
3. A polynucleotide as claimed in claim 1,  
wherein the polynucleotide is an RNA.
4. A polynucleotide as claimed in claim 1,  
30 comprising the nucleotide sequence as shown in SEQ ID  
No. 4 or SEQ ID No. 9.

5. A DNA as claimed in claim 2 which is capable of replication, comprising
- (i) one or more nucleotide sequences shown in SEQ ID No. 4 or SEQ ID No. 9, or
  - 5 (ii) at least one sequence which corresponds to sequence (i) within the range of the degeneration of the genetic code, or
  - (iii) at least one sequence which hybridizes with the sequence complementary to sequence (i) or (ii),  
10 and optionally
  - (iv) sense mutations of neutral function in (i).
6. A polynucleotide as claimed in claim 1, which codes for a polypeptide which comprises at least one of the amino acid sequences as shown in SEQ ID  
15 No. 3, SEQ ID No. 5, SEQ ID NO. 8 and SEQ ID No. 10.
7. A coryneform microorganism, in particular of the genus Corynebacterium, transformed by the introduction of the DNA, which is capable of replication, as claimed in one of claims 1 or 5.
- 20 8. A process for the preparation of L-amino acids, which comprises carrying out the following steps,
- a) fermentation of the bacteria which produce the desired L-amino acid, in which preferably at  
25 least the zwf gene and optionally one or more of the genes tkt gene or devB gene are amplified, in addition to the opcA gene.
  - b) concentration of the desired product in the medium or in the cells of the bacteria and cells  
30 of the bacteria concentrated and
  - c) isolation of the desired L-amino acid.

9. The process as claimed in claim 8,  
w h e r e i n  
one or more further gene(s) of the pentose phosphate  
cycle is or are amplified, in particular over-  
5 expressed, in addition to the genes mentioned.
10. The process as claimed in claims 8 and 9,  
w h e r e i n  
to achieve the amplification, the number of copies of  
the genes or nucleotide sequences is increased by  
10 transformation of the microorganisms with plasmid  
vectors which carry these genes or nucleotide  
sequences.
11. The process as claimed in claim 8,  
w h e r e i n  
15 for the preparation of amino acids, in particular  
lysine, bacteria in which, in addition to the opca  
gene, one or more genes chosen from the group  
consisting of
- 20 11.1 the dapA gene which codes for  
dihydrodipicolinate synthase,  
11.2 the lysC gene which codes for a feed back  
resistant aspartate kinase,  
11.3 the gap gene which codes for glycerolaldehyde 3-  
25 phosphate dehydrogenase,  
11.4 the pyc gene which codes for pyruvate  
carboxylase,  
11.5 the tkt gene which codes for transketolase,  
11.6 the gnd gene which codes for 6-phosphogluconate  
30 dehydrogenase,  
11.7 the lysE gene which codes for lysine export,  
11.8 the zwf gene,  
11.9 the eno gene which codes for enolase,  
11.10 the tal gene which codes for transaldolase,  
35 11.11 in particular the zwf gene

is or are amplified, in particular over-expressed, at the same time, are fermented.

12. The process as claimed in claim 8,  
w h e r e i n

5 for the preparation of amino acids, in particular L-lysine, bacteria in which one or more genes chosen from the group consisting of

12.1 the pck gene which codes for phosphoenol pyruvate  
10 carboxykinase,

12.2 the pgi gene which codes for glucose 6-phosphate  
isomerase,

12.3 the poxB gene which codes for pyruvate oxidase or

12.4 the zwa2 gene

15 is or are attenuated at the same time, are fermented.

13. The use of polynucleotide sequences as claimed in claim 1 as primers for the preparation of DNA of genes which display an effect corresponding to the opcA gene by the polymerase chain reaction.

20 14. The use of polynucleotide sequences as claimed in claim 1 as hybridization probes.




Figure 1:




Figure 2:






Figure 3:

## SEQUENCE PROTOCOL

<110> National University of Ireland, Galway  
 Degussa-Hüls AG  
 5       Forschungszentrum Juelich GmbH  
  
 <120> New nucleotide sequences which code for the opcA gene  
  
 <130> 990213 BT  
 10  
 <140>  
 <141>  
  
 <160> 12  
 15  
 <170> PatentIn Ver. 2.1  
  
 <210> 1  
 <211> 6995  
 20 <212> DNA  
 <213> Corynebacterium glutamicum ATCC13032  
  
 <220>  
 <221> CDS  
 25 <222> (3658)..(5202)  
 <223> zwf  
  
 <220>  
 <221> CDS  
 30 <222> (5217)..(6173)  
 <223> opcA  
  
 <400> 1  
 35 cacatttgaa ccacagttgg ttataaaatg ggttcaacat cactatgggt agaggtggtg 60  
 acgggtcaga ttaagcaaag actactttcg gggtagatca cctttgcaa atttgaacca 120  
 attaacctaa gtcgtagatc tgatcatcgg atctaacgaa aacgaaccaa aactttggtc 180  
 40 ccggtttaac ccaggaagga ttgaccaact tgacgctgtc acctgaactt caggcgctca 240  
 ctgtacgcaa ttaccctctt gattggtccg atgtggacac caaggctgta gacactgttc 300  
 gtgtcctcgc tgcagacgct gtagaaaact gtggctccgg ccaccaggc accgcaatga 360  
 45 gcctgggtcc ccttgcatat accttgtagc agcgggttat gaacgtagat ccacaggaca 420  
 ccaactgggc aggcctgac cgcttcgttc tttcttggtg cactcctct ttgaccagtc 480  
 50 acatccagct ttacttgggt ggattcggcc ttgagatgga tgacctgaag gctctgcgca 540  
 cctgggattc cttgacccca ggacacctg agtaccgcca caccaagggc gttgagatca 600  
 ccactggccc tcttgccag ggtcttgcat ctgcagttgg tatggccatg gctgctcgtc 660  
 55 gtgagcgtgg cctattcgac ccaaccgctg ctgagggcga atccccattc gaccaccaca 720  
 tctacgtcat tgcttctgat ggtgacctgc aggaaggtgt cacctctgag gcactctcca 780  
 60 tgcgtggcac ccagcagctg ggcaacctca tcgtgttctg ggatgacaac cgcactctcca 840  
 tcgaagacaa cactgagatc gctttcaacg aggacgttgt tgctcgttac aaggcttacg 900

gctggcagac cattgaggtt gaggctggcg aggacgttgc agcaatcgaa gctgcagtgg 960  
ctgaggctaa gaaggacacc aagcgaccta ccttcacccg cgttcgcacc atcatcggct 1020  
5 tcccagctcc aactatgatg aacaccgggtg ctgtgcacgg tgctgctctt ggcgcagctg 1080  
aggttgcagc aaccaagact gagcttggat tcgacccga ggctcacttc gcgatcgacg 1140  
atgaggttat cgctcacacc cgctccctcg cagagcgcg cgcacagaag aaggctgcat 1200  
10 ggcaggtcaa gtctgatgag tgggcagctg ccaaccctga gaacaaggct ctgttcgac 1260  
gcctgaactc ccgtgagctt ccagcgggct acgctgacga gctcccaaca tgggatgcag 1320  
15 atgagaaggg cgctcgcaact cgtaaggctt ccgaggctgc acttcaggca ctgggcaaga 1380  
cccttcctga gctgtggggc gggtccgctg acctcgagg ttccaacaac accgtgatca 1440  
agggctcccc ttccttcggc cctgagtcca tctccaccga gacctggtct gctgagcctt 1500  
20 acggccgtaa cctgcacttc ggtatccgtg agcacgctat gggatccatc ctcaacggca 1560  
tttccctcca cgggtggcacc cgccatacgc gcggaacctt cctcatcttc tccgactaca 1620  
25 tgctgctgc agttcgtctt gcagctctca tggagaccga cgcttactac gtctggacc 1680  
acgactccat cggctcgggc gaagatggcc caaccacca gcctgttgaa accttggtg 1740  
cactgcgcgc catcccaggt ctgtccgtcc tgctgctgc agatgcgaac gagaccgcc 1800  
30 aggcttgggc tgcagcactt gagtacaagg aaggccctaa ggtcttgca ctgaccgcc 1860  
agaacgttcc tgttctggaa ggcaccaagg agaaggctgc tgaaggcgtt cgccgcggtg 1920  
35 gctacgtcct ggttgagggt tccaaggaaa cccagatgt gatcctcatg ggctccggct 1980  
ccgaggttca gcttgagctt aacgctgcga aggtctgga agctgagggc gttgcagctc 2040  
gcgttgcttc cgttccttgc atggattggt tccaggagca ggacgcagag tacatcgagt 2100  
40 ccgttctgcc tgcagctgtg accgctcgtg tgtctgtga agctggcatc gcaatgcctt 2160  
ggtaaccgtt cttgggcacc cagggccgtg ctgtctccct tgagcacttc ggtgcttctg 2220  
45 cggattacca gaccctgttt gagaagttcg gcatcaccac cgatgcagtc gtggcagcgg 2280  
ccaaggactc cattaacggt taattgccct gctgttttta gcttcaacc ggggcaatat 2340  
gattctccgg aattttattg ccccggttg ttgtgttaa tcggtacaaa ggtcttaag 2400  
50 cacatccctt acttgctgc tctccttgag cacagttcaa gaacaattct tttaaggaaa 2460  
atttagtttc atgtctcaca ttgatgatct tgcacagctc ggcacttcca cttggctcga 2520  
55 cgacctctcc cgcgagcgca ttactccgg caatctcagc caggttattg agggaaagtc 2580  
tgtagtcggt gtcaccacca acccagctat tttcgagca gcaatgtcca agggcgattc 2640  
ctacgacgct cagatcgag agctcaaggc cgctggcgca tctgttgacc aggtgttta 2700  
60 cgccatgagc atcgacgacg ttcgcaatgc ttgtgatctg ttcaccggca tcttcgagtc 2760  
ctccaacggc tacgacggcc gcgtgtccat cgaggttgac ccacgtatct ctgctgaccg 2820

cgacgcaacc ctggctcagg ccaaggagct gtgggcaaag gttgatcgtc caaacgtcat 2880  
 gatcaagatc cctgcaaccc caggttcttt gccagcaatc accgacgctt tggctgaggg 2940  
 5 catcagcggt aacgtcacct tgatcttctc cgttgctcgc taccgcgagg tcacgcgtgc 3000  
 gttcatcgag ggcatcaagc aggtgctgc aaacggccac gacgtctcca agatccactc 3060  
 10 tgtggcttcc ttcttcgtct cccgcgtcga cgttgagatc gacaagcgcc tcgaggcaat 3120  
 cggatccgat gaggctttgg ctctgcgcgg caaggcaggc gttgccaacg ctcagcgcg 3180  
 ttacgctgtg tacaaggagc ttttcgacgc cgccgagctg cctgaagggtg ccaacactca 3240  
 15 gcgcccactg tgggcatcca ccggcgtgaa gaacctgcg tacgctgcaa ctctttacgt 3300  
 ttccgagctg gctggtccaa acaccgtcaa caccatgcca gaaggcacca tcgacgcggt 3360  
 20 tctggagcag ggcaacctgc acggtgacac cctgtccaac tccgcggcag aagctgacgc 3420  
 tgtgttctcc cagcttgagg ctctgggctg tgacttgcca gatgtcttcc aggtcctgga 3480  
 gaccgaggggt gtggacaagt tcgttgcttc ttggagcgaa ctgcttgagt ccatggaagc 3540  
 25 tcgcctgaag tagaatcagc acgtgcac acgtacggcg acatgaaatc gaattagttc 3600  
 gatcttatgt ggccgttaca catctttcat taaagaaagg atcgtgacac taccatc 3657  
 30 gtg agc aca aac acg acc ccc tcc agc tgg aca aac cca ctg cgc gac 3705  
 Val Ser Thr Asn Thr Thr Pro Ser Ser Trp Thr Asn Pro Leu Arg Asp  
 1 5 10 15  
 ccg cag gat aaa cga ctc ccc cgc atc gct ggc cct tcc ggc atg gtg 3753  
 35 Pro Gln Asp Lys Arg Leu Pro Arg Ile Ala Gly Pro Ser Gly Met Val  
 20 25 30  
 atc ttc ggt gtc act ggc gac ttg gct cga aag aag ctg ctc ccc gcc 3801  
 40 Ile Phe Gly Val Thr Gly Asp Leu Ala Arg Lys Lys Leu Leu Pro Ala  
 35 40 45  
 att tat gat cta gca aac cgc gga ttg ctg ccc cca gga ttc tcg ttg 3849  
 Ile Tyr Asp Leu Ala Asn Arg Gly Leu Leu Pro Pro Gly Phe Ser Leu  
 50 55 60  
 45 gta ggt tac ggc cgc cgc gaa tgg tcc aaa gaa gac ttt gaa aaa tac 3897  
 Val Gly Tyr Gly Arg Arg Glu Trp Ser Lys Glu Asp Phe Glu Lys Tyr  
 65 70 75 80

	gta cgc gat gcc gca agt gct ggt gct cgt acg gaa ttc cgt gaa aat	3945
	Val Arg Asp Ala Ala Ser Ala Gly Ala Arg Thr Glu Phe Arg Glu Asn	
	85 90 95	
5	ggt tgg gag cgc ctc gcc gag ggt atg gaa ttt gtt cgc ggc aac ttt	3993
	Val Trp Glu Arg Leu Ala Glu Gly Met Glu Phe Val Arg Gly Asn Phe	
	100 105 110	
10	gat gat gat gca gct ttc gac aac ctc gct gca aca ctc aag cgc atc	4041
	Asp Asp Asp Ala Ala Phe Asp Asn Leu Ala Ala Thr Leu Lys Arg Ile	
	115 120 125	
15	gac aaa acc cgc ggc acc gcc ggc aac tgg gct tac tac ctg tcc att	4089
	Asp Lys Thr Arg Gly Thr Ala Gly Asn Trp Ala Tyr Tyr Leu Ser Ile	
	130 135 140	
20	cca cca gat tcc ttc aca gcg gtc tgc cac cag ctg gag cgt tcc ggc	4137
	Pro Pro Asp Ser Phe Thr Ala Val Cys His Gln Leu Glu Arg Ser Gly	
	145 150 155 160	
	atg gct gaa tcc acc gaa gaa gca tgg cgc cgc gtg atc atc gag aag	4185
	Met Ala Glu Ser Thr Glu Glu Ala Trp Arg Arg Val Ile Ile Glu Lys	
	165 170 175	
25	cct ttc ggc cac aac ctc gaa tcc gca cac gag ctc aac cag ctg gtc	4233
	Pro Phe Gly His Asn Leu Glu Ser Ala His Glu Leu Asn Gln Leu Val	
	180 185 190	
30	aac gca gtc ttc cca gaa tct tct gtg ttc cgc atc gac cac tat ttg	4281
	Asn Ala Val Phe Pro Glu Ser Ser Val Phe Arg Ile Asp His Tyr Leu	
	195 200 205	
35	ggc aag gaa aca gtt caa aac atc ctg gct ctg cgt ttt gct aac cag	4329
	Gly Lys Glu Thr Val Gln Asn Ile Leu Ala Leu Arg Phe Ala Asn Gln	
	210 215 220	
40	ctg ttt gag cca ctg tgg aac tcc aac tac gtt gac cac gtc cag atc	4377
	Leu Phe Glu Pro Leu Trp Asn Ser Asn Tyr Val Asp His Val Gln Ile	
	225 230 235 240	
	acc atg gct gaa gat att ggc ttg ggt gga cgt gct ggt tac tac gac	4425
	Thr Met Ala Glu Asp Ile Gly Leu Gly Gly Arg Ala Gly Tyr Tyr Asp	
	245 250 255	
45	ggc atc ggc gca gcc cgc gac gtc atc cag aac cac ctg atc cag ctc	4473
	Gly Ile Gly Ala Ala Arg Asp Val Ile Gln Asn His Leu Ile Gln Leu	
	260 265 270	
50	ttg gct ctg gtt gcc atg gaa gaa cca att tct ttc gtg cca gcg cag	4521
	Leu Ala Leu Val Ala Met Glu Glu Pro Ile Ser Phe Val Pro Ala Gln	
	275 280 285	
55	ctg cag gca gaa aag atc aag gtg ctc tct gcg aca aag ccg tgc tac	4569
	Leu Gln Ala Glu Lys Ile Lys Val Leu Ser Ala Thr Lys Pro Cys Tyr	
	290 295 300	

	cca ttg gat aaa acc tcc gct cgt ggt cag tac gct gcc ggt tgg cag	4617
	Pro Leu Asp Lys Thr Ser Ala Arg Gly Gln Tyr Ala Ala Gly Trp Gln	
	305 310 315 320	
5	ggc tct gag tta gtc aag gga ctt cgc gaa gaa gat ggc ttc aac cct	4665
	Gly Ser Glu Leu Val Lys Gly Leu Arg Glu Glu Asp Gly Phe Asn Pro	
	325 330 335	
10	gag tcc acc act gag act ttt gcg gct tgt acc tta gag atc acg tct	4713
	Glu Ser Thr Thr Glu Thr Phe Ala Ala Cys Thr Leu Glu Ile Thr Ser	
	340 345 350	
15	cgt cgc tgg gct ggt gtg ccg ttc tac ctg cgc acc ggt aag cgt ctt	4761
	Arg Arg Trp Ala Gly Val Pro Phe Tyr Leu Arg Thr Gly Lys Arg Leu	
	355 360 365	
20	ggt cgc cgt gtt act gag att gcc gtg gtg ttt aaa gac gca cca cac	4809
	Gly Arg Arg Val Thr Glu Ile Ala Val Val Phe Lys Asp Ala Pro His	
	370 375 380	
25	cag cct ttc gac ggc gac atg act gta tcc ctt ggc caa aac gcc atc	4857
	Gln Pro Phe Asp Gly Asp Met Thr Val Ser Leu Gly Gln Asn Ala Ile	
	385 390 395 400	
	gtg att cgc gtg cag cct gat gaa ggt gtg ctc atc cgc ttc ggt tcc	4905
	Val Ile Arg Val Gln Pro Asp Glu Gly Val Leu Ile Arg Phe Gly Ser	
	405 410 415	
30	aag gtt cca ggt tct gcc atg gaa gtc cgt gac gtc aac atg gac ttc	4953
	Lys Val Pro Gly Ser Ala Met Glu Val Arg Asp Val Asn Met Asp Phe	
	420 425 430	
35	tcc tac tca gaa tcc ttc act gaa gaa tca cct gaa gca tac gag cgc	5001
	Ser Tyr Ser Glu Ser Phe Thr Glu Glu Ser Pro Glu Ala Tyr Glu Arg	
	435 440 445	
40	ctc att ttg gat gcg ctg tta gat gaa tcc agc ctc ttc cct acc aac	5049
	Leu Ile Leu Asp Ala Leu Leu Asp Glu Ser Ser Leu Phe Pro Thr Asn	
	450 455 460	
45	gag gaa gtg gaa ctg agc tgg aag att ctg gat cca att ctt gaa gca	5097
	Glu Glu Val Glu Leu Ser Trp Lys Ile Leu Asp Pro Ile Leu Glu Ala	
	465 470 475 480	
	tgg gat gcc gat gga gaa cca gag gat tac cca gcg ggt acg tgg ggt	5145
	Trp Asp Ala Asp Gly Glu Pro Glu Asp Tyr Pro Ala Gly Thr Trp Gly	
	485 490 495	
50	cca aag agc gct gat gaa atg ctt tcc cgc aac ggt cac acc tgg cgc	5193
	Pro Lys Ser Ala Asp Glu Met Leu Ser Arg Asn Gly His Thr Trp Arg	
	500 505 510	
55	agg cca taa tttaggggca aaaa atg atc ttt gaa ctt ccg gat acc acc	5243
	Arg Pro 515 Met Ile Phe Glu Leu Pro Asp Thr Thr	
	520	

	acc	cag	caa	att	tcc	aag	acc	cta	act	cga	ctg	cgt	gaa	tcg	ggc	acc	5291
	Thr	Gln	Gln	Ile	Ser	Lys	Thr	Leu	Thr	Arg	Leu	Arg	Glu	Ser	Gly	Thr	
	525					530					535					540	
5	cag	gtc	acc	acc	ggc	cga	gtg	ctc	acc	ctc	atc	gtg	gtc	act	gac	tcc	5339
	Gln	Val	Thr	Thr	Gly	Arg	Val	Leu	Thr	Leu	Ile	Val	Val	Thr	Asp	Ser	
					545					550					555		
10	gaa	agc	gat	gtc	gct	gca	gtt	acc	gag	tcc	acc	aat	gaa	gcc	tcg	cgc	5387
	Glu	Ser	Asp	Val	Ala	Ala	Val	Thr	Glu	Ser	Thr	Asn	Glu	Ala	Ser	Arg	
				560					565					570			
15	gag	cac	cca	tct	cgc	gtg	atc	att	ttg	gtg	gtt	ggc	gat	aaa	act	gca	5435
	Glu	His	Pro	Ser	Arg	Val	Ile	Ile	Leu	Val	Val	Gly	Asp	Lys	Thr	Ala	
			575					580					585				
20	gaa	aac	aaa	gtt	gac	gca	gaa	gtc	cgt	atc	ggg	ggc	gac	gct	ggg	gct	5483
	Glu	Asn	Lys	Val	Asp	Ala	Glu	Val	Arg	Ile	Gly	Gly	Asp	Ala	Gly	Ala	
		590					595					600					
25	tcc	gag	atg	atc	atc	atg	cat	ctc	aac	gga	cct	gtc	gct	gac	aag	ctc	5531
	Ser	Glu	Met	Ile	Ile	Met	His	Leu	Asn	Gly	Pro	Val	Ala	Asp	Lys	Leu	
	605					610					615					620	
	cag	tat	gtc	gtc	aca	cca	ctg	ttg	ctt	cct	gac	acc	ccc	atc	gtt	gct	5579
	Gln	Tyr	Val	Val	Thr	Pro	Leu	Leu	Leu	Pro	Asp	Thr	Pro	Ile	Val	Ala	
					625					630					635		
30	tgg	tgg	cca	ggg	gaa	tca	cca	aag	aat	cct	tcc	cag	gac	cca	att	gga	5627
	Trp	Trp	Pro	Gly	Glu	Ser	Pro	Lys	Asn	Pro	Ser	Gln	Asp	Pro	Ile	Gly	
				640				645						650			
35	cgc	atc	gca	caa	cga	cgc	atc	act	gat	gct	ttg	tac	gac	cgt	gat	gac	5675
	Arg	Ile	Ala	Gln	Arg	Arg	Ile	Thr	Asp	Ala	Leu	Tyr	Asp	Arg	Asp	Asp	
			655					660					665				
40	gca	cta	gaa	gat	cgt	gtt	gag	aac	tat	cac	cca	ggg	gat	acc	gac	atg	5723
	Ala	Leu	Glu	Asp	Arg	Val	Glu	Asn	Tyr	His	Pro	Gly	Asp	Thr	Asp	Met	
		670					675					680					
45	acg	tgg	gcg	cgc	ctt	acc	cag	tgg	cgg	gga	ctt	gtt	gcc	tcc	tca	ttg	5771
	Thr	Trp	Ala	Arg	Leu	Thr	Gln	Trp	Arg	Gly	Leu	Val	Ala	Ser	Ser	Leu	
	685					690					695					700	
	gat	cac	cca	cca	cac	agc	gaa	atc	act	tcc	gtg	agg	ctg	acc	ggg	gca	5819
	Asp	His	Pro	Pro	His	Ser	Glu	Ile	Thr	Ser	Val	Arg	Leu	Thr	Gly	Ala	
					705					710					715		
50	agc	ggc	agt	acc	tcg	gtg	gat	ttg	gct	gca	ggc	tgg	ttg	gcg	cgg	agg	5867
	Ser	Gly	Ser	Thr	Ser	Val	Asp	Leu	Ala	Ala	Gly	Trp	Leu	Ala	Arg	Arg	
				720				725						730			
55	ctg	aaa	gtg	cct	gtg	atc	cgc	gag	gtg	aca	gat	gct	ccc	acc	gtg	cca	5915
	Leu	Lys	Val	Pro	Val	Ile	Arg	Glu	Val	Thr	Asp	Ala	Pro	Thr	Val	Pro	
			735					740					745				

acc gat gag ttt ggt act cca ctg ctg gct atc cag cgc ctg gag atc 5963  
 Thr Asp Glu Phe Gly Thr Pro Leu Leu Ala Ile Gln Arg Leu Glu Ile  
 750 755 760

5 gtt cgc acc acc ggc tcg atc atc atc acc atc tat gac gct cat acc 6011  
 Val Arg Thr Thr Gly Ser Ile Ile Ile Thr Ile Tyr Asp Ala His Thr  
 765 770 775 780

10 ctt cag gta gag atg ccg gaa tcc ggc aat gcc cca tcg ctg gtg gct 6059  
 Leu Gln Val Glu Met Pro Glu Ser Gly Asn Ala Pro Ser Leu Val Ala  
 785 790 795

15 att ggt cgt cga agt gag tcc gac tgc ttg tct gag gag ctt cgc cac 6107  
 Ile Gly Arg Arg Ser Glu Ser Asp Cys Leu Ser Glu Glu Leu Arg His  
 800 805 810

20 atg gat cca gat ttg ggc tac cag cac gca cta tcc ggc ttg tcc agc 6155  
 Met Asp Pro Asp Leu Gly Tyr Gln His Ala Leu Ser Gly Leu Ser Ser  
 815 820 825

gtc aag ctg gaa acc gtc taaggagaaa tacaacacta tgggtgatgt 6203  
 Val Lys Leu Glu Thr Val  
 830

25 agtacgcgca cgcgatactg aagatttgggt tgcacaggct gcctccaaat tcattgaggt 6263  
 tgttgaagca gcaactgccca ataattggcac cgcacaggta gtgctcaccg gtggtggcgc 6323

30 cggcatcaag ttgctggaag agctcagcgt tgatgcggct gaccttgctt gggatcgcat 6383  
 tcatgtgttc ttccggcgatg agcgcaatgt ccctgtcagt gattctgagt ccaatgaggg 6443  
 ccaggctcgt gaggcactgt tgtccaaggt ttctatccct gaagccaaca ttcacggata 6503

35 tgggtctcggc gacgtagatc ttgcagaggc agcccgcgct tacgaagctg tgttgatga 6563  
 attcgcacca aacggctttg atcttcacct gctcggcatg ggtggcgaag gccatatcaa 6623

40 ctccctgttc cctcacaccg atgcagtcaa ggaatcctcc gaaagggtca tcgcggtgtt 6683  
 tgattcccct aagcctcctt cagagcgtgc aactctaacc cttcctgcgg ttactccgc 6743  
 aaagcgcgtg tgggtgctgg tttctggtgc ggagaaggct gaggcagctg cggcgatcgt 6803

45 caacggtgag cctgctgttg agtggcctgc tgctggagct accggatctg aggaaacggt 6863  
 attgttcttg gctgatgatg ctgcaggaaa tctctaagca gcgccagctc taacaagaag 6923

50 ctttaacaag aagctctaac gaaaagcact aacaaactaa tccgggtgcg aaccttcac 6983  
 tgaatcgatg ga 6995

55 <210> 2  
 <211> 514  
 <212> PRT  
 <213> Corynebacterium glutamicum ATCC13032

60 <400> 2  
 Val Ser Thr Asn Thr Thr Pro Ser Ser Trp Thr Asn Pro Leu Arg Asp  
 1 5 10 15



Pro Gln Asp Lys Arg Leu Pro Arg Ile Ala Gly Pro Ser Gly Met Val  
                   20                  25                  30  
 5 Ile Phe Gly Val Thr Gly Asp Leu Ala Arg Lys Lys Leu Leu Pro Ala  
                   35                  40                  45  
 Ile Tyr Asp Leu Ala Asn Arg Gly Leu Leu Pro Pro Gly Phe Ser Leu  
                   50                  55                  60  
 10 Val Gly Tyr Gly Arg Arg Glu Trp Ser Lys Glu Asp Phe Glu Lys Tyr  
                   65                  70                  75                  80  
 Val Arg Asp Ala Ala Ser Ala Gly Ala Arg Thr Glu Phe Arg Glu Asn  
                   85                  90                  95  
 15 Val Trp Glu Arg Leu Ala Glu Gly Met Glu Phe Val Arg Gly Asn Phe  
                   100                  105  
 Asp Asp Asp Ala Ala Phe Asp Asn Leu Ala Ala Thr Leu Lys Arg Ile  
                   115                  120                  125  
 Asp Lys Thr Arg Gly Thr Ala Gly Asn Trp Ala Tyr Tyr Leu Ser Ile  
                   130                  135                  140  
 25 Pro Pro Asp Ser Phe Thr Ala Val Cys His Gln Leu Glu Arg Ser Gly  
                   145                  150                  155                  160  
 Met Ala Glu Ser Thr Glu Glu Ala Trp Arg Arg Val Ile Ile Glu Lys  
                   165                  170                  175  
 30 Pro Phe Gly His Asn Leu Glu Ser Ala His Glu Leu Asn Gln Leu Val  
                   180                  185                  190  
 Asn Ala Val Phe Pro Glu Ser Ser Val Phe Arg Ile Asp His Tyr Leu  
                   195                  200                  205  
 Gly Lys Glu Thr Val Gln Asn Ile Leu Ala Leu Arg Phe Ala Asn Gln  
                   210                  215                  220  
 40 Leu Phe Glu Pro Leu Trp Asn Ser Asn Tyr Val Asp His Val Gln Ile  
                   225                  230                  235                  240  
 Thr Met Ala Glu Asp Ile Gly Leu Gly Gly Arg Ala Gly Tyr Tyr Asp  
                   245                  250                  255  
 45 Gly Ile Gly Ala Ala Arg Asp Val Ile Gln Asn His Leu Ile Gln Leu  
                   260                  265                  270  
 Leu Ala Leu Val Ala Met Glu Glu Pro Ile Ser Phe Val Pro Ala Gln  
                   275                  280                  285  
 50 Leu Gln Ala Glu Lys Ile Lys Val Leu Ser Ala Thr Lys Pro Cys Tyr  
                   290                  295                  300

Pro Leu Asp Lys Thr Ser Ala Arg Gly Gln Tyr Ala Ala Gly Trp Gln  
 305 310 315 320  
 5 Gly Ser Glu Leu Val Lys Gly Leu Arg Glu Glu Asp Gly Phe Asn Pro  
 325 330 335  
 Glu Ser Thr Thr Glu Thr Phe Ala Ala Cys Thr Leu Glu Ile Thr Ser  
 340 345 350  
 10 Arg Arg Trp Ala Gly Val Pro Phe Tyr Leu Arg Thr Gly Lys Arg Leu  
 355 360 365  
 15 Gly Arg Arg Val Thr Glu Ile Ala Val Val Phe Lys Asp Ala Pro His  
 370 375 380  
 Gln Pro Phe Asp Gly Asp Met Thr Val Ser Leu Gly Gln Asn Ala Ile  
 385 390 395 400  
 20 Val Ile Arg Val Gln Pro Asp Glu Gly Val Leu Ile Arg Phe Gly Ser  
 405 410 415  
 Lys Val Pro Gly Ser Ala Met Glu Val Arg Asp Val Asn Met Asp Phe  
 420 425 430  
 25 Ser Tyr Ser Glu Ser Phe Thr Glu Glu Ser Pro Glu Ala Tyr Glu Arg  
 435 440 445  
 30 Leu Ile Leu Asp Ala Leu Leu Asp Glu Ser Ser Leu Phe Pro Thr Asn  
 450 455 460  
 Glu Glu Val Glu Leu Ser Trp Lys Ile Leu Asp Pro Ile Leu Glu Ala  
 465 470 475 480  
 35 Trp Asp Ala Asp Gly Glu Pro Glu Asp Tyr Pro Ala Gly Thr Trp Gly  
 485 490 495  
 Pro Lys Ser Ala Asp Glu Met Leu Ser Arg Asn Gly His Thr Trp Arg  
 500 505 510  
 40 Arg Pro  
 <210> 3  
 45 <211> 319  
 <212> PRT  
 <213> Corynebacterium glutamicum ATCC13032  
 <400> 3  
 50 Met Ile Phe Glu Leu Pro Asp Thr Thr Thr Gln Gln Ile Ser Lys Thr  
 1 5 10 15  
 Leu Thr Arg Leu Arg Glu Ser Gly Thr Gln Val Thr Thr Gly Arg Val  
 20 25 30  
 55 Leu Thr Leu Ile Val Val Thr Asp Ser Glu Ser Asp Val Ala Ala Val  
 35 40 45  
 60 Thr Glu Ser Thr Asn Glu Ala Ser Arg Glu His Pro Ser Arg Val Ile  
 50 55 60  
 Ile Leu Val Val Gly Asp Lys Thr Ala Glu Asn Lys Val Asp Ala Glu

	65		70		75		80
	Val Arg Ile Gly Gly Asp Ala Gly Ala Ser Glu Met Ile Ile Met His						
		85			90		95
5	Leu Asn Gly Pro Val Ala Asp Lys Leu Gln Tyr Val Val Thr Pro Leu						
		100			105		110
10	Leu Leu Pro Asp Thr Pro Ile Val Ala Trp Trp Pro Gly Glu Ser Pro						
		115			120		125
	Lys Asn Pro Ser Gln Asp Pro Ile Gly Arg Ile Ala Gln Arg Arg Ile						
		130			135		140
15	Thr Asp Ala Leu Tyr Asp Arg Asp Asp Ala Leu Glu Asp Arg Val Glu						
		145			150		155
	Asn Tyr His Pro Gly Asp Thr Asp Met Thr Trp Ala Arg Leu Thr Gln						
		165			170		175
20	Trp Arg Gly Leu Val Ala Ser Ser Leu Asp His Pro Pro His Ser Glu						
		180			185		190
	Ile Thr Ser Val Arg Leu Thr Gly Ala Ser Gly Ser Thr Ser Val Asp						
25		195			200		205
	Leu Ala Ala Gly Trp Leu Ala Arg Arg Leu Lys Val Pro Val Ile Arg						
		210			215		220
30	Glu Val Thr Asp Ala Pro Thr Val Pro Thr Asp Glu Phe Gly Thr Pro						
		225			230		235
	Leu Leu Ala Ile Gln Arg Leu Glu Ile Val Arg Thr Thr Gly Ser Ile						
		245			250		255
35	Ile Ile Thr Ile Tyr Asp Ala His Thr Leu Gln Val Glu Met Pro Glu						
		260			265		270
	Ser Gly Asn Ala Pro Ser Leu Val Ala Ile Gly Arg Arg Ser Glu Ser						
40		275			280		285
	Asp Cys Leu Ser Glu Glu Leu Arg His Met Asp Pro Asp Leu Gly Tyr						
		290			295		300
45	Gln His Ala Leu Ser Gly Leu Ser Ser Val Lys Leu Glu Thr Val						
		305			310		315
50	<210> 4						
	<211> 960						
	<212> DNA						
	<213> Corynebacterium glutamicum ATCC13032						
55	<220>						
	<221> CDS						
	<222> (1)..(957)						
	<223> opcA						
60	<400> 4						
	atg atc ttt gaa ctt ccg gat acc acc acc cag caa att tcc aag acc						48
	Met Ile Phe Glu Leu Pro Asp Thr Thr Thr Gln Gln Ile Ser Lys Thr						
	1	5			10		15

11/23

	cta act cga ctg cgt gaa tcg ggc acc cag gtc acc acc ggc cga gtg	96
	Leu Thr Arg Leu Arg Glu Ser Gly Thr Gln Val Thr Thr Gly Arg Val	
	20 25 30	
5	ctc acc ctc atc gtg gtc act gac tcc gaa agc gat gtc gct gca gtt	144
	Leu Thr Leu Ile Val Val Thr Asp Ser Glu Ser Asp Val Ala Ala Val	
	35 40 45	
10	acc gag tcc acc aat gaa gcc tcg cgc gag cac cca tct cgc gtg atc	192
	Thr Glu Ser Thr Asn Glu Ala Ser Arg Glu His Pro Ser Arg Val Ile	
	50 55 60	
15	att ttg gtg gtt ggc gat aaa act gca gaa aac aaa gtt gac gca gaa	240
	Ile Leu Val Val Gly Asp Lys Thr Ala Glu Asn Lys Val Asp Ala Glu	
	65 70 75 80	
20	gtc cgt atc ggt ggc gac gct ggt gct tcc gag atg atc atc atg cat	288
	Val Arg Ile Gly Gly Asp Ala Gly Ala Ser Glu Met Ile Ile Met His	
	85 90 95	
25	ctc aac gga cct gtc gct gac aag ctc cag tat gtc gtc aca cca ctg	336
	Leu Asn Gly Pro Val Ala Asp Lys Leu Gln Tyr Val Val Thr Pro Leu	
	100 105 110	
30	ttg ctt cct gac acc ccc atc gtt gct tgg tgg cca ggt gaa tca cca	384
	Leu Leu Pro Asp Thr Pro Ile Val Ala Trp Trp Pro Gly Glu Ser Pro	
	115 120 125	
35	aag aat cct tcc cag gac cca att gga cgc atc gca caa cga cgc atc	432
	Lys Asn Pro Ser Gln Asp Pro Ile Gly Arg Ile Ala Gln Arg Arg Ile	
	130 135 140	
40	act gat gct ttg tac gac cgt gat gac gca cta gaa gat cgt gtt gag	480
	Thr Asp Ala Leu Tyr Asp Arg Asp Asp Ala Leu Glu Asp Arg Val Glu	
	145 150 155 160	
45	aac tat cac cca ggt gat acc gac atg acg tgg gcg cgc ctt acc cag	528
	Asn Tyr His Pro Gly Asp Thr Asp Met Thr Trp Ala Arg Leu Thr Gln	
	165 170 175	
50	tgg cgg gga ctt gtt gcc tcc tca ttg gat cac cca cca cac agc gaa	576
	Trp Arg Gly Leu Val Ala Ser Ser Leu Asp His Pro Pro His Ser Glu	
	180 185 190	
55	atc act tcc gtg agg ctg acc ggt gca agc ggc agt acc tcg gtg gat	624
	Ile Thr Ser Val Arg Leu Thr Gly Ala Ser Gly Ser Thr Ser Val Asp	
	195 200 205	



Thr Asp Ala Leu Tyr Asp Arg Asp Asp Ala Leu Glu Asp Arg Val Glu  
 145 150 155 160  
 5 Asn Tyr His Pro Gly Asp Thr Asp Met Thr Trp Ala Arg Leu Thr Gln  
 165 170 175  
 Trp Arg Gly Leu Val Ala Ser Ser Leu Asp His Pro Pro His Ser Glu  
 180 185 190  
 10 Ile Thr Ser Val Arg Leu Thr Gly Ala Ser Gly Ser Thr Ser Val Asp  
 195 200 205  
 Leu Ala Ala Gly Trp Leu Ala Arg Arg Leu Lys Val Pro Val Ile Arg  
 210 215 220  
 15 Glu Val Thr Asp Ala Pro Thr Val Pro Thr Asp Glu Phe Gly Thr Pro  
 225 230 235 240  
 20 Leu Leu Ala Ile Gln Arg Leu Glu Ile Val Arg Thr Thr Gly Ser Ile  
 245 250 255  
 Ile Ile Thr Ile Tyr Asp Ala His Thr Leu Gln Val Glu Met Pro Glu  
 260 265 270  
 25 Ser Gly Asn Ala Pro Ser Leu Val Ala Ile Gly Arg Arg Ser Glu Ser  
 275 280 285  
 Asp Cys Leu Ser Glu Glu Leu Arg His Met Asp Pro Asp Leu Gly Tyr  
 290 295 300  
 30 Gln His Ala Leu Ser Gly Leu Ser Ser Val Lys Leu Glu Thr Val  
 305 310 315  
 35 <210> 6  
 <211> 3038  
 <212> DNA  
 <213> Corynebacterium glutamicum AS019  
 40 <220>  
 <221> CDS  
 <222> (115)..(1659)  
 <223> zwf  
 45 <220>  
 <221> CDS  
 <222> (1672)..(2628)  
 <223> opcA  
 50 <400> 6  
 cctgaagtag aatcagcacg ctgcatcagt aacggcgaca tgaaatcgaa ttagttcgat 60

	cttatgtggc cgttacacat ctttcattaa agaaaggatc gtgacactac catc gtg	117
	Val	1
5	agc aca aac acg acc ccc tcc agc tgg aca aac cca ctg cgc gac ccg	165
	Ser Thr Asn Thr Pro Ser Ser Trp Thr Asn Pro Leu Arg Asp Pro	5 10 15
10	cag gat aaa cga ctc ccc cgc atc gct ggc cct tcc ggc atg gtg atc	213
	Gln Asp Lys Arg Leu Pro Arg Ile Ala Gly Pro Ser Gly Met Val Ile	20 25 30
15	ttc ggt gtc act ggc gac ttg gct cga aag aag ctg ctc ccc gcc att	261
	Phe Gly Val Thr Gly Asp Leu Ala Arg Lys Lys Leu Leu Pro Ala Ile	35 40 45
20	tat gat cta gca aac cgc gga ttg ctg ccc cca gga ttc tcg ttg gta	309
	Tyr Asp Leu Ala Asn Arg Gly Leu Leu Pro Pro Gly Phe Ser Leu Val	50 55 60 65
25	ggt tac ggc cgc cgc gaa tgg tcc aaa gaa gac ttt gaa aaa tac gta	357
	Gly Tyr Gly Arg Arg Glu Trp Ser Lys Glu Asp Phe Glu Lys Tyr Val	70 75 80
30	cgc gat gcc gca agt gct ggt gct cgt acg gaa ttc cgt gaa aat gtt	405
	Arg Asp Ala Ala Ser Ala Gly Ala Arg Thr Glu Phe Arg Glu Asn Val	85 90 95
35	tgg gag cgc ctc gcc gag ggt atg gaa ttt gtt cgc ggc aac ttt gat	453
	Trp Glu Arg Leu Ala Glu Gly Met Glu Phe Val Arg Gly Asn Phe Asp	100 105 110
40	gat gat gca gct ttc gac aac ctc gct gca aca ctc aag cgc atc gac	501
	Asp Asp Ala Ala Phe Asp Asn Leu Ala Ala Thr Leu Lys Arg Ile Asp	115 120 125
45	aaa acc cgc ggc acc gcc ggc aac tgg gct tac tac ctg tcc att cca	549
	Lys Thr Arg Gly Thr Ala Gly Asn Trp Ala Tyr Tyr Leu Ser Ile Pro	130 135 140 145
50	cca gat tcc ttc aca gcg gtc tgc cac cag ctg gag cgt tcc ggc atg	597
	Pro Asp Ser Phe Thr Ala Val Cys His Gln Leu Glu Arg Ser Gly Met	150 155 160
55	gct gaa tcc acc gaa gaa gca tgg cgc cgc gtg atc atc gag aag cct	645
	Ala Glu Ser Thr Glu Glu Ala Trp Arg Arg Val Ile Ile Glu Lys Pro	165 170 175
60	ttc ggc cac aac ctc gaa tcc gca cac gag ctc aac cag ctg gtc aac	693
	Phe Gly His Asn Leu Glu Ser Ala His Glu Leu Asn Gln Leu Val Asn	180 185 190
65	gca gtc ttc cca gaa tct tct gtg ttc cgc atc gac cac tat ttg ggc	741
	Ala Val Phe Pro Glu Ser Ser Val Phe Arg Ile Asp His Tyr Leu Gly	195 200 205
70	aag gaa aca gtt caa aac atc ctg gct ctg cgt ttt gct aac cag ctg	789
	Lys Glu Thr Val Gln Asn Ile Leu Ala Leu Arg Phe Ala Asn Gln Leu	210 215 220 225
75	ttt gag cca ctg tgg aac tcc aac tac gtt gac cac gtc cag atc acc	837
	Phe Glu Pro Leu Trp Asn Ser Asn Tyr Val Asp His Val Gln Ile Thr	230 235 240

	atg gct gaa gat att ggc ttg ggt gga cgt gct ggt tac tac gac ggc	885
	Met Ala Glu Asp Ile Gly Leu Gly Gly Arg Ala Gly Tyr Tyr Asp Gly	
	245 250 255	
5	atc ggc gca ccg cgc gac gtc atc cag aac cac ctg atc cag ctc ttg	933
	Ile Gly Ala Pro Arg Asp Val Ile Gln Asn His Leu Ile Gln Leu Leu	
	260 265 270	
10	gct ctg gtt gcc atg gaa gaa cca att tct ttc gtg cca gcg gca cgg	981
	Ala Leu Val Ala Met Glu Glu Pro Ile Ser Phe Val Pro Ala Ala Arg	
	275 280 285	
15	cag gca gaa aag atc aag gtg ctc tct gcg aca aag ccg tgc tac cca	1029
	Gln Ala Glu Lys Ile Lys Val Leu Ser Ala Thr Lys Pro Cys Tyr Pro	
	290 295 300 305	
20	ttg gat aaa acc tcc gct cgt ggt cag tac gct gcc ggt tgg cag ggc	1077
	Leu Asp Lys Thr Ser Ala Arg Gly Gln Tyr Ala Ala Gly Trp Gln Gly	
	310 315 320	
25	tct gag tta gtc aag gga ctt cgc gaa gaa gat ggc ttc aac cct gag	1125
	Ser Glu Leu Val Lys Gly Leu Arg Glu Glu Asp Gly Phe Asn Pro Glu	
	325 330 335	
30	tcc acc act gag act ttt gcg gct tgt acc tta gag atc acg tct cgt	1173
	Ser Thr Thr Glu Thr Phe Ala Ala Cys Thr Leu Glu Ile Thr Ser Arg	
	340 345 350	
35	cgc tgg gct ggt gtg ccg ttc tac ctg cgc acc ggt aag cgt ctt ggt	1221
	Arg Trp Ala Gly Val Pro Phe Tyr Leu Arg Thr Gly Lys Arg Leu Gly	
	355 360 365	
40	cgc cgt gtt act gag att gcc gtg gtg ttt aaa gac gca cca cac cag	1269
	Arg Arg Val Thr Glu Ile Ala Val Val Phe Lys Asp Ala Pro His Gln	
	370 375 380 385	
45	cct ttc gac ggc gac atg act gta tcc ctt ggc caa aac gcc atc gtg	1317
	Pro Phe Asp Gly Asp Met Thr Val Ser Leu Gly Gln Asn Ala Ile Val	
	390 395 400	
50	att cgc gtg cag cct gat gaa ggt gtg ctc atc cgc ttc ggt tcc aag	1365
	Ile Arg Val Gln Pro Asp Glu Gly Val Leu Ile Arg Phe Gly Ser Lys	
	405 410 415	
50	gtt cca ggt tct gcc atg gaa gtc cgt gac gtc aac atg gac ttc tcc	1413
	Val Pro Gly Ser Ala Met Glu Val Arg Asp Val Asn Met Asp Phe Ser	
	420 425 430	
50	tac tca gaa tcc ttc act gaa gaa tca cct gaa gca tac gag cgc ctc	1461
	Tyr Ser Glu Ser Phe Thr Glu Glu Ser Pro Glu Ala Tyr Glu Arg Leu	
	435 440 445	



	att ttg gat gcg ctg tta gat gaa tcc agc ctc ttc cct acc aac gag	1509
	Ile Leu Asp Ala Leu Leu Asp Glu Ser Ser Leu Phe Pro Thr Asn Glu	
	450 455 460 465	
5		
	gaa gtg gaa ctg agc tgg aag att ctg gat cca att ctt gaa gca tgg	1557
	Glu Val Glu Leu Ser Trp Lys Ile Leu Asp Pro Ile Leu Glu Ala Trp	
	470 475 480	
10		
	gat gcc gat gga gaa cca gag gat tac cca gcg ggt acg tgg ggt cca	1605
	Asp Ala Asp Gly Glu Pro Glu Asp Tyr Pro Ala Gly Thr Trp Gly Pro	
	485 490 495	
15		
	aag agc gct gat gaa atg ctt tcc cgc aac ggt cac acc tgg cgc agg	1653
	Lys Ser Ala Asp Glu Met Leu Ser Arg Asn Gly His Thr Trp Arg Arg	
	500 505 510	
20		
	cca taa tttaggggca aa atg atc ttt gaa ctt ccg gat acc acc acc cag	1704
	Pro Met Ile Phe Glu Leu Pro Asp Thr Thr Gln	
	515 520 525	
25		
	caa att tcc aag acc cta act cga ctg cgt gaa tcg ggc acc cag gtc	1752
	Gln Ile Ser Lys Thr Leu Thr Arg Leu Arg Glu Ser Gly Thr Gln Val	
	530 535 540	
30		
	acc acc ggc cga gtg ctc acc ctc atc gtg gtc act gac tcc gaa agc	1800
	Thr Thr Gly Arg Val Leu Thr Leu Ile Val Val Thr Asp Ser Glu Ser	
	545 550 555	
35		
	gat gtc gct gca gtt acc gag tcc acc aat gaa gcc tcg cgc gag cac	1848
	Asp Val Ala Ala Val Thr Glu Ser Thr Asn Glu Ala Ser Arg Glu His	
	560 565 570	
40		
	cca tct cgc gtg atc att ttg gtg gtt ggc gat aaa act gca gaa aac	1896
	Pro Ser Arg Val Ile Ile Leu Val Val Gly Asp Lys Thr Ala Glu Asn	
	575 580 585 590	
45		
	aaa gtt gac gca gaa gtc cgt atc ggt ggc gac gct ggt gct tcc gag	1944
	Lys Val Asp Ala Glu Val Arg Ile Gly Gly Asp Ala Gly Ala Ser Glu	
	595 600 605	
50		
	atg atc atc atg cat ctc aac gga cct gtc gct gac aag ctc cag tat	1992
	Met Ile Ile Met His Leu Asn Gly Pro Val Ala Asp Lys Leu Gln Tyr	
	610 615 620	
55		
	gtc gtc aca cca ctg ttg ctt cct gac acc ccc atc gtt gct tgg tgg	2040
	Val Val Thr Pro Leu Leu Leu Pro Asp Thr Pro Ile Val Ala Trp Trp	
	625 630 635	
60		
	cca ggt gaa tca cca aag aat cct tcc cag gac cca att gga cgc atc	2088
	Pro Gly Glu Ser Pro Lys Asn Pro Ser Gln Asp Pro Ile Gly Arg Ile	
	640 645 650	
65		
	gca caa cga cgc atc act gat gct ttg tac gac cgt gat gac gca cta	2136
	Ala Gln Arg Arg Ile Thr Asp Ala Leu Tyr Asp Arg Asp Ala Leu	
	655 660 665 670	

	gaa gat cgt gtt gag aac tat cac cca ggt gat acc gac atg acg tgg	2184
	Glu Asp Arg Val Glu Asn Tyr His Pro Gly Asp Thr Asp Met Thr Trp	
	675 680 685	
5	gcg cgc ctt acc cag tgg cgg gga ctt gtt gcc tcc tca ttg gat cac	2232
	Ala Arg Leu Thr Gln Trp Arg Gly Leu Val Ala Ser Ser Leu Asp His	
	690 695 700	
10	cca cca cac agc gaa atc act tcc gtg agg ctg acc ggt gca agc ggc	2280
	Pro Pro His Ser Glu Ile Thr Ser Val Arg Leu Thr Gly Ala Ser Gly	
	705 710 715	
15	agt acc tcg gtg gat ttg gct gca ggc tgg ttg gcg cgg agg ctg aaa	2328
	Ser Thr Ser Val Asp Leu Ala Ala Gly Trp Leu Ala Arg Arg Leu Lys	
	720 725 730	
20	gtg cct gtg atc cgc gag gtg aca gat gct ccc acc gtg cca acc gat	2376
	Val Pro Val Ile Arg Glu Val Thr Asp Ala Pro Thr Val Pro Thr Asp	
	735 740 745 750	
	gag ttt ggt act cca ctg ctg gct atc cag cgc ctg gag atc gtt cgc	2424
	Glu Phe Gly Thr Pro Leu Leu Ala Ile Gln Arg Leu Glu Ile Val Arg	
	755 760 765	
25	acc acc ggc tcg atc atc atc acc atc tat gac gct cat acc ctt cag	2472
	Thr Thr Gly Ser Ile Ile Ile Thr Ile Tyr Asp Ala His Thr Leu Gln	
	770 775 780	
30	gta gag atg ccg gaa tcc ggc aat gcc cca tcg ctg gtg gct att ggt	2520
	Val Glu Met Pro Glu Ser Gly Asn Ala Pro Ser Leu Val Ala Ile Gly	
	785 790 795	
35	cgt cga agt gag tcc gac tgc ttg tct gag gag ctt cgc cac atg gat	2568
	Arg Arg Ser Glu Ser Asp Cys Leu Ser Glu Glu Leu Arg His Met Asp	
	800 805 810	
40	cca gat ttg ggc tac cag cac gca cta tcc ggc ttg tcc agc gtc aag	2616
	Pro Asp Leu Gly Tyr Gln His Ala Leu Ser Gly Leu Ser Ser Val Lys	
	815 820 825 830	
	ctg gaa acc gtc taaggagaaa tacaacacta tggttgatgt agtacgcgca	2668
	Leu Glu Thr Val	
45	cgcatactga agatttgggt gcacaggctg cctccaaatt cattgagggt gttgaagcag	2728
	caactgccaa taatggcacc gcacaggtag tgctcaccgg ttgtggcgcc ggcatcaagt	2788
	tgctggaaaa gctcagcgtt gatgcggctg accttcctg ggatcgcatt catgtgttct	2848
50	tcggcgatga gcgcaatgtc cctgtcagtg attctgagtc caatgagggc caggctcgtg	2908
	aggcactgtt gtccaagggt tctatccctg aagccaacat tcacggatat ggtctcggcg	2968
55	acgtagatct tgcagaggca gcccgcgctt acgaagctgt gttggatgaa ttcgcaccaa	3028
	acggccttga	3038
60	<210> 7	
	<211> 514	
	<212> PRT	
	<213> Corynebacterium glutamicum AS019	

<400> 7  
 Val Ser Thr Asn Thr Thr Pro Ser Ser Trp Thr Asn Pro Leu Arg Asp  
 1 5 10 15  
 5 Pro Gln Asp Lys Arg Leu Pro Arg Ile Ala Gly Pro Ser Gly Met Val  
 20 25 30  
 10 Ile Phe Gly Val Thr Gly Asp Leu Ala Arg Lys Lys Leu Leu Pro Ala  
 35 40 45  
 Ile Tyr Asp Leu Ala Asn Arg Gly Leu Leu Pro Pro Gly Phe Ser Leu  
 50 55 60  
 15 Val Gly Tyr Gly Arg Arg Glu Trp Ser Lys Glu Asp Phe Glu Lys Tyr  
 65 70 75 80  
 Val Arg Asp Ala Ala Ser Ala Gly Ala Arg Thr Glu Phe Arg Glu Asn  
 85 90 95  
 20 Val Trp Glu Arg Leu Ala Glu Gly Met Glu Phe Val Arg Gly Asn Phe  
 100 105 110  
 25 Asp Asp Asp Ala Ala Phe Asp Asn Leu Ala Ala Thr Leu Lys Arg Ile  
 115 120 125  
 Asp Lys Thr Arg Gly Thr Ala Gly Asn Trp Ala Tyr Tyr Leu Ser Ile  
 130 135 140  
 30 Pro Pro Asp Ser Phe Thr Ala Val Cys His Gln Leu Glu Arg Ser Gly  
 145 150 155 160  
 Met Ala Glu Ser Thr Glu Glu Ala Trp Arg Arg Val Ile Ile Glu Lys  
 165 170 175  
 35 Pro Phe Gly His Asn Leu Glu Ser Ala His Glu Leu Asn Gln Leu Val  
 180 185 190  
 40 Asn Ala Val Phe Pro Glu Ser Ser Val Phe Arg Ile Asp His Tyr Leu  
 195 200 205  
 Gly Lys Glu Thr Val Gln Asn Ile Leu Ala Leu Arg Phe Ala Asn Gln  
 210 215 220  
 45 Leu Phe Glu Pro Leu Trp Asn Ser Asn Tyr Val Asp His Val Gln Ile  
 225 230 235 240  
 Thr Met Ala Glu Asp Ile Gly Leu Gly Gly Arg Ala Gly Tyr Tyr Asp  
 245 250 255  
 50 Gly Ile Gly Ala Pro Arg Asp Val Ile Gln Asn His Leu Ile Gln Leu  
 260 265 270  
 55 Leu Ala Leu Val Ala Met Glu Glu Pro Ile Ser Phe Val Pro Ala Ala  
 275 280 285  
 Arg Gln Ala Glu Lys Ile Lys Val Leu Ser Ala Thr Lys Pro Cys Tyr  
 290 295 300  
 60 Pro Leu Asp Lys Thr Ser Ala Arg Gly Gln Tyr Ala Ala Gly Trp Gln  
 305 310 315 320  
 Gly Ser Glu Leu Val Lys Gly Leu Arg Glu Glu Asp Gly Phe Asn Pro

325 330 335  
 Glu Ser Thr Thr Glu Thr Phe Ala Ala Cys Thr Leu Glu Ile Thr Ser  
 340 345 350  
 5 Arg Arg Trp Ala Gly Val Pro Phe Tyr Leu Arg Thr Gly Lys Arg Leu  
 355 360 365  
 10 Gly Arg Arg Val Thr Glu Ile Ala Val Val Phe Lys Asp Ala Pro His  
 370 375 380  
 Gln Pro Phe Asp Gly Asp Met Thr Val Ser Leu Gly Gln Asn Ala Ile  
 385 390 395 400  
 15 Val Ile Arg Val Gln Pro Asp Glu Gly Val Leu Ile Arg Phe Gly Ser  
 405 410 415  
 Lys Val Pro Gly Ser Ala Met Glu Val Arg Asp Val Asn Met Asp Phe  
 420 425 430  
 20 Ser Tyr Ser Glu Ser Phe Thr Glu Glu Ser Pro Glu Ala Tyr Glu Arg  
 435 440 445  
 25 Leu Ile Leu Asp Ala Leu Leu Asp Glu Ser Ser Leu Phe Pro Thr Asn  
 450 455 460  
 Glu Glu Val Glu Leu Ser Trp Lys Ile Leu Asp Pro Ile Leu Glu Ala  
 465 470 475 480  
 30 Trp Asp Ala Asp Gly Glu Pro Glu Asp Tyr Pro Ala Gly Thr Trp Gly  
 485 490 495  
 Pro Lys Ser Ala Asp Glu Met Leu Ser Arg Asn Gly His Thr Trp Arg  
 500 505 510  
 35 Arg Pro  
 <210> 8  
 40 <211> 319  
 <212> PRT  
 <213> Corynebacterium glutamicum AS019  
 <400> 8  
 45 Met Ile Phe Glu Leu Pro Asp Thr Thr Thr Gln Gln Ile Ser Lys Thr  
 1 5 10 15  
 Leu Thr Arg Leu Arg Glu Ser Gly Thr Gln Val Thr Thr Gly Arg Val  
 20 25 30

Leu Thr Leu Ile Val Val Thr Asp Ser Glu Ser Asp Val Ala Ala Val  
 35 40 45  
 5 Thr Glu Ser Thr Asn Glu Ala Ser Arg Glu His Pro Ser Arg Val Ile  
 50 55 60  
 Ile Leu Val Val Gly Asp Lys Thr Ala Glu Asn Lys Val Asp Ala Glu  
 65 70 75 80  
 10 Val Arg Ile Gly Gly Asp Ala Gly Ala Ser Glu Met Ile Ile Met His  
 85 90 95  
 Leu Asn Gly Pro Val Ala Asp Lys Leu Gln Tyr Val Val Thr Pro Leu  
 100 105 110  
 Leu Leu Pro Asp Thr Pro Ile Val Ala Trp Trp Pro Gly Glu Ser Pro  
 115 120 125  
 20 Lys Asn Pro Ser Gln Asp Pro Ile Gly Arg Ile Ala Gln Arg Arg Ile  
 130 135 140  
 Thr Asp Ala Leu Tyr Asp Arg Asp Asp Ala Leu Glu Asp Arg Val Glu  
 145 150 155 160  
 25 Asn Tyr His Pro Gly Asp Thr Asp Met Thr Trp Ala Arg Leu Thr Gln  
 165 170 175  
 Trp Arg Gly Leu Val Ala Ser Ser Leu Asp His Pro Pro His Ser Glu  
 180 185 190  
 Ile Thr Ser Val Arg Leu Thr Gly Ala Ser Gly Ser Thr Ser Val Asp  
 195 200 205  
 35 Leu Ala Ala Gly Trp Leu Ala Arg Arg Leu Lys Val Pro Val Ile Arg  
 210 215 220  
 Glu Val Thr Asp Ala Pro Thr Val Pro Thr Asp Glu Phe Gly Thr Pro  
 225 230 235 240  
 40 Leu Leu Ala Ile Gln Arg Leu Glu Ile Val Arg Thr Thr Gly Ser Ile  
 245 250 255  
 Ile Ile Thr Ile Tyr Asp Ala His Thr Leu Gln Val Glu Met Pro Glu  
 260 265 270  
 45 Ser Gly Asn Ala Pro Ser Leu Val Ala Ile Gly Arg Arg Ser Glu Ser  
 275 280 285  
 50 Asp Cys Leu Ser Glu Glu Leu Arg His Met Asp Pro Asp Leu Gly Tyr  
 290 295 300  
 Gln His Ala Leu Ser Gly Leu Ser Ser Val Lys Leu Glu Thr Val  
 305 310 315  
 55

&lt;210&gt; 9

&lt;211&gt; 960

60 &lt;212&gt; DNA

&lt;213&gt; Corynebacterium glutamicum AS019

&lt;220&gt;

<221> CDS  
 <222> (1)..(957)  
 <223> opca

```

5  <400> 9
   atg atc ttt gaa ctt ccg gat acc acc acc cag caa att tcc aag acc 48
   Met Ile Phe Glu Leu Pro Asp Thr Thr Thr Gln Gln Ile Ser Lys Thr
   1                               10                               15

10  cta act cga ctg cgt gaa tcg ggc acc cag gtc acc acc ggc cga gtg 96
   Leu Thr Arg Leu Arg Glu Ser Gly Thr Thr Gln Val Thr Thr Gly Arg Val
   20                               25                               30

15  ctc acc ctc atc gtg gtc act gac tcc gaa agc gat gtc gct gca gtt 144
   Leu Thr Leu Ile Val Val Thr Asp Ser Glu Ser Asp Val Ala Ala Val
   35                               40                               45

20  acc gag tcc acc aat gaa gcc tcg cgc gag cac cca tct cgc gtg atc 192
   Thr Glu Ser Thr Asn Glu Ala Ser Arg Glu His Pro Ser Arg Val Ile
   50                               55                               60

25  att ttg gtg gtt ggc gat aaa act gca gaa aac aaa gtt gac gca gaa 240
   Ile Leu Val Val Gly Asp Lys Thr Ala Glu Asn Lys Val Asp Ala Glu
   65                               70                               75                               80

   gtc cgt atc ggt ggc gac gct ggt gct tcc gag atg atc atc atg cat 288
   Val Arg Ile Gly Gly Asp Ala Gly Ala Ser Glu Met Ile Ile Met His
   85                               90                               95

30  ctc aac gga cct gtc gct gac aag ctc cag tat gtc gtc aca cca ctg 336
   Leu Asn Gly Pro Val Ala Asp Lys Leu Gln Tyr Val Val Thr Pro Leu
   100                               105                               110

35  ttg ctt cct gac acc ccc atc gtt gct tgg tgg cca ggt gaa tca cca 384
   Leu Leu Pro Asp Thr Pro Ile Val Ala Trp Trp Pro Gly Glu Ser Pro
   115                               120                               125

40  aag aat cct tcc cag gac cca att gga cgc atc gca caa cga cgc atc 432
   Lys Asn Pro Ser Gln Asp Pro Ile Gly Arg Ile Ala Gln Arg Arg Ile
   130                               135                               140

   act gat gct ttg tac gac cgt gat gac gca cta gaa gat cgt gtt gag 480
   Thr Asp Ala Leu Tyr Asp Arg Asp Asp Ala Leu Glu Asp Arg Val Glu
   145                               150                               155                               160

45  aac tat cac cca ggt gat acc gac atg acg tgg gcg cgc ctt acc cag 528
   Asn Tyr His Pro Gly Asp Thr Asp Met Thr Trp Ala Arg Leu Thr Gln
   165                               170                               175

50  tgg cgg gga ctt gtt gcc tcc tca ttg gat cac cca cca cac agc gaa 576
   Trp Arg Gly Leu Val Ala Ser Ser Leu Asp His Pro Pro His Ser Glu
   180                               185                               190

```

	atc act tcc gtg agg ctg acc ggt gca agc ggc agt acc tcg gtg gat	624
	Ile Thr Ser Val Arg Leu Thr Gly Ala Ser Gly Ser Thr Ser Val Asp	
	195 200 205	
5	ttg gct gca ggc tgg ttg gcg cgg agg ctg aaa gtg cct gtg atc cgc	672
	Leu Ala Ala Gly Trp Leu Ala Arg Arg Leu Lys Val Pro Val Ile Arg	
	210 215 220	
10	gag gtg aca gat gct ccc acc gtg cca acc gat gag ttt ggt act cca	720
	Glu Val Thr Asp Ala Pro Thr Val Pro Thr Asp Glu Phe Gly Thr Pro	
	225 230 235 240	
15	ctg ctg gct atc cag cgc ctg gag atc gtt cgc acc acc ggc tcg atc	768
	Leu Leu Ala Ile Gln Arg Leu Glu Ile Val Arg Thr Thr Gly Ser Ile	
	245 250 255	
20	atc atc acc atc tat gac gct cat acc ctt cag gta gag atg ccg gaa	816
	Ile Ile Thr Ile Tyr Asp Ala His Thr Leu Gln Val Glu Met Pro Glu	
	260 265 270	
25	tcc ggc aat gcc cca tcg ctg gtg gct att ggt cgt cga agt gag tcc	864
	Ser Gly Asn Ala Pro Ser Leu Val Ala Ile Gly Arg Arg Ser Glu Ser	
	275 280 285	
30	gac tgc ttg tct gag gag ctt cgc cac atg gat cca gat ttg ggc tac	912
	Asp Cys Leu Ser Glu Glu Leu Arg His Met Asp Pro Asp Leu Gly Tyr	
	290 295 300	
35	cag cac gca cta tcc ggc ttg tcc agc gtc aag ctg gaa acc gtc taa	960
	Gln His Ala Leu Ser Gly Leu Ser Ser Val Lys Leu Glu Thr Val	
	305 310 315	
40	<210> 10	
	<211> 319	
	<212> PRT	
	<213> Corynebacterium glutamicum AS019	
45	<400> 10	
	Met Ile Phe Glu Leu Pro Asp Thr Thr Thr Gln Gln Ile Ser Lys Thr	
	1 5 10 15	
50	Leu Thr Arg Leu Arg Glu Ser Gly Thr Gln Val Thr Thr Gly Arg Val	
	20 25 30	
55	Leu Thr Leu Ile Val Val Thr Asp Ser Glu Ser Asp Val Ala Ala Val	
	35 40 45	
60	Thr Glu Ser Thr Asn Glu Ala Ser Arg Glu His Pro Ser Arg Val Ile	
	50 55 60	
65	Ile Leu Val Val Gly Asp Lys Thr Ala Glu Asn Lys Val Asp Ala Glu	
	65 70 75 80	
70	Val Arg Ile Gly Gly Asp Ala Gly Ala Ser Glu Met Ile Ile Met His	
	85 90 95	
75	Leu Asn Gly Pro Val Ala Asp Lys Leu Gln Tyr Val Val Thr Pro Leu	
	100 105 110	
80	Leu Leu Pro Asp Thr Pro Ile Val Ala Trp Trp Pro Gly Glu Ser Pro	
	115 120 125	

Lys Asn Pro Ser Gln Asp Pro Ile Gly Arg Ile Ala Gln Arg Arg Ile  
 130 135 140  
 5 Thr Asp Ala Leu Tyr Asp Arg Asp Asp Ala Leu Glu Asp Arg Val Glu  
 145 150 155 160  
 Asn Tyr His Pro Gly Asp Thr Asp Met Thr Trp Ala Arg Leu Thr Gln  
 165 170 175  
 10 Trp Arg Gly Leu Val Ala Ser Ser Leu Asp His Pro Pro His Ser Glu  
 180 185 190  
 Ile Thr Ser Val Arg Leu Thr Gly Ala Ser Gly Ser Thr Ser Val Asp  
 195 200 205  
 15 Leu Ala Ala Gly Trp Leu Ala Arg Arg Leu Lys Val Pro Val Ile Arg  
 210 215 220  
 20 Glu Val Thr Asp Ala Pro Thr Val Pro Thr Asp Glu Phe Gly Thr Pro  
 225 230 235 240  
 Leu Leu Ala Ile Gln Arg Leu Glu Ile Val Arg Thr Thr Gly Ser Ile  
 245 250 255  
 25 Ile Ile Thr Ile Tyr Asp Ala His Thr Leu Gln Val Glu Met Pro Glu  
 260 265 270  
 Ser Gly Asn Ala Pro Ser Leu Val Ala Ile Gly Arg Arg Ser Glu Ser  
 275 280 285  
 30 Asp Cys Leu Ser Glu Glu Leu Arg His Met Asp Pro Asp Leu Gly Tyr  
 290 295 300  
 35 Gln His Ala Leu Ser Gly Leu Ser Ser Val Lys Leu Glu Thr Val  
 305 310 315  
 40 <210> 11  
 <211> 15  
 <212> PRT  
 <213> Corynebacterium glutamicum ATCC13032  
 <400> 11  
 45 Xaa Xaa Xaa Xaa Xaa Pro Xaa Xaa Trp Xaa Asn Pro Leu Arg Asp  
 1 5 10 15  
 50 <210> 12  
 <211> 15  
 <212> PRT  
 <213> Corynebacterium glutamicum ATCC13032  
 55 <400> 12  
 Met Ile Phe Xaa Leu Pro Asp Xaa Xaa Xaa Gln Gln Ile Ser Lys  
 1 5 10 15  
 60



## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 00/06300

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/53 C12N9/04 C12P13/08 C12N1/21 C12Q1/68  
/(C12N1/21,C12R1:15),(C12P13/08,C12R1:15)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C12P C12Q C12R

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, EMBL, WPI Data, BIOSIS, MEDLINE, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EMBL 'Online! Database Entry E13655 , 23 June 1998 (1998-06-23) HATAKEYAMA K. ET AL.: "gDNA encoding glucose-6-phosphate dehydrogenase" Database accession no. E13655 XP002152311	1,2,5, 13,14
Y	the whole document	8
X	& JP 09 224661 A (MITSUBISHI CHEM CORP) 2 September 1997 (1997-09-02) cited in the application the whole document	1,2,5,7, 8,13,14
	— —/—	

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*A\* document member of the same patent family

Date of the actual completion of the international search

9 November 2000

Date of mailing of the international search report

22/11/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5018 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3010

Authorized officer

Montero Lopez, B

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 00/06300

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>DATABASE WPI Section Ch, Week 198824 Derwent Publications Ltd., London, GB; Class B10, AN 1988-164726 XP002152313</p> <p>KATSUMATA, RYOICHI ET AL.: "L-Glutamic acid production by culturing <i>Corynebacterium</i> or <i>Brevibacterium</i> transformed with recombinant DNA containing information for production of phospho:fructo:kinase" &amp; JP 63 102692 A (KYOWA HAKKO KOGYO KK), 7 May 1988 (1988-05-07) abstract</p>	8
A	<p>YOSHINORI KOBAYASHI ET AL.: "Purification and properties of NAD-dependent D-Glucose dehydrogenase produced by alkalophilic <i>Corynebacterium</i> sp. No. 93-1" AGRICULTURAL AND BIOLOGICAL CHEMISTRY, vol. 44, no. 10, 1980, pages 2261-2269, XP002152310 TOKYO JP the whole document</p>	1-14
A	<p>DATABASE EMBL 'Online! Database Entry SP33285, 11 October 1995 (1995-10-11) NEWMAN J. ET AL.: "Synechococcus PCC7942 zwf region, fructose 1,6-biphosphatase (fbp), glucose 6-phosphate dehydrogenase (zwf), OpcA (opcA), cytochrome b6 (petD), and cytochrome b6f complex subunit IV (petB) genes, complete cds." Database accession no. U33285 XP002152312 the whole document &amp; JULIE NEWMAN ET AL.: "A comparison of gene organization in the zwf region of the genomes of the cyanobacteria <i>Synechococcus</i> sp. PCC 7942 and <i>Anabaena</i> sp. PCC 7120" FEMS MICROBIOLOGY LETTERS, vol. 133, no. 1-2, 1995, pages 187-193, XP000957662</p>	1-14

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 00/06300

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>MICHAEL L. SUMMERS ET AL.: "Transcriptional regulation of zwf, encoding glucose-6-phosphate dehydrogenase, from the cyanobacterium Nostoc punctiforme strain ATCC 29133" MOLECULAR MICROBIOLOGY, vol. 22, no. 3, 1996, pages 473-480, XP000961316 summary page 477, left-hand column, paragraph 3 -page 479, left-hand column, paragraph 3</p>	1-7

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 00/06300

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
JP 9224661 A	02-09-1997	NONE	
JP 63102692 A	07-05-1988	JP 2101579 C JP 7121227 B	22-10-1996 25-12-1995